

BANKS, H.

AM
1920
ban

THE RELATIVE SIGNIFICANCE
of
CARBONIC ACID AND AMMONIA ELIMINATION
in the
ACID INTOXICATIONS OF PREGNANCY.

* * * *

Bibliographical thesis
presented for partial credit
toward the A.M. degree.

* * * *

by
HELEN LAMBERT BANKS, B.S.
BOSTON UNIVERSITY, 1918.

B78.744
B6
1920 AM
Gann

Digitized by the Internet Archive
in 2014

<https://archive.org/details/relativesignific00bank>

Introduction.

I. Study of the history of the subject.
with the view of ascertaining
the state of knowledge.

II. Scope.

1. Definition of the subject.
2. Limits of the subject.
3. Scope of the subject.

III. Method.

1. General method.
2. Special method.
3. Details of the method.
4. Results of the method.

Summary of the main results.

I. Classification and terminology.

Summary of the main results.

I. Classification and terminology.

Conclusion.

I. Criteria for selection of material.

II. Methods chosen.

III. Plan of research.

Bibliography.

P1127



1515

O U T L I N E .

Introduction.

I. Functional Disturbances Normally Associated with the State of Pregnancy.

(a) Acid-Intoxications.

1. Symptoms.

(a) Elimination of Acetone-bodies.

(b) Lessening of the Titratable

Alkalinity of the Blood.

2. Etiology.

3. Clinical Determinations Available for the Estimation of Approaching, and Measurement of an Existing Acid-intoxication.

Methods of Ammonia Determination.

I. Classification and Criticism.

Methods of Alveolar Carbon Dioxide Determinations.

I. Classification and Criticism.

Conclusion.

I. Criteria for Selection of Methods.

II. Methods Chosen.

III. Plan of Research.

Bibliography.

OUTLINE.

Introduction.

I. Functional Disturbances Normally Associated

with the State of Pregnancy.

(a) Acid-Intoxications.

I. Symptoms.

(a) Elimination of Acetone-bodies.

(b) Lessening of the Titratable

Alkalinity of the Blood.

2. Etiology.

3. Clinical Determinations Available for

the Estimation of Approaching, and

Measurement of an Existing Acid-intox-

ication.

Methods of Ammonia Determination.

I. Classification and Criticism.

Methods of Alveolar Carbon Dioxide Determinations.

I. Classification and Criticism.

Conclusion.

I. Criteria for Selection of Methods.

II. Methods Chosen.

III. Plan of Research.

Bibliography.

INTRODUCTION

THE RELATIVE SIGNIFICANCE OF CARBONIC ACID
AND AMMONIA ELIMINATION
IN THE ACID INTOXICATIONS OF PREGNANCY.

* * * * *

Of the various normal functions of the animal body the state of pregnancy as an incident of the reproductive function shows a curious anomaly. That the reproductive functions are to be regarded as fundamental and, hence, as wholly normal in character is too obvious to require discussion.

That, on the other hand, during the state of pregnancy, certain disturbances of normal function form characteristic incidents of the condition, is a general and well-recognized fact. The severity of these functional disturbances varies profoundly with the individual, so that while in many instances they are so superficial as to be wholly without significance, in an appreciable percentage they assume a gravity of character which may seriously endanger not only the health, but even the life of both mother and child. The general picture of these anomalies is the same, the differences being those of degree and not of kind.

Among these functional disturbances may be classed the various toxemias, several different types of which are recognized as potentially incidental to the pregnant state. Of these the so-called acid intoxications form a relatively

THE RELATIVE SIGNIFICANCE OF CARBONIC ACID

AND AMMONIA ELIMINATION

IN THE ACID INTOXICATION OF PREGNANCY.

* * * * *

Of the various normal functions of the animal

body the state of pregnancy as an incident of the reproductive function shows a curious anomaly. That the reproductive functions are to be regarded as fundamental and, hence, as wholly normal in character is too obvious to require discussion.

That, on the other hand, during the state of pregnancy, certain disturbances of normal function form characteristic incidents of the condition, is a general and well-recognized fact. The severity of these functional disturbances varies profoundly with the individual, so that while in many instances they are so superficial as to be wholly without significance, in an appreciable percentage they assume a gravity of character which may seriously endanger not only the health, but even the life of both mother and child. The general picture of these anomalies is the same, the differences being those of degree and not of kind.

Among these functional disturbances may be classed the various toxemias, several different types of which are recognized as potentially incidental to the pregnant state. Of these the so-called acid intoxications form a relatively

common and highly important group. It is necessary here, again, to differentiate between two types of acid intoxication. In the first of these, an imperfect oxidation of one or more of the normal energy-producing food materials may be regarded as the basis of the so-called acidosis. It is distinguished by a typical group of symptoms, among which may be mentioned the elimination of the so-called acetone bodies - B-hydroxybutyric acid, acetoacetic acid, and acetone - by the kidneys, a lessening of the titratable alkalinity of the blood, the hydrogen ion content remaining practically unchanged, a diminution of the carbon dioxide content of the alveolar air and a marked increase of the ammonia content of the urine. This group of symptoms is further found in certain other pathological and abnormal physiological states. For example, it is found that the feeding of mineral acids to herbivorous animals will produce a lessening of the titratable alkalinity of the blood, a diminution in the alveolar carbonic acid, and an increase in the urine of the fixed bases of the animal body, as sodium, potassium, calcium, and magnesium. The generally accepted explanation of this phenomenon, is that the acid on passing into the blood stream unites with the alkaline elements of the blood, and through the very delicate regulation of salt concentrations conditioned by functional activity in the kidney, there is a definite loss of these alkaline elements, the normal function of which is to fix the carbon dioxide thrown off during the katabolic activity of the cell, and thus permit

common and highly important group. It is necessary here, again, to differentiate between two types of acid intoxication. In the first of these, an imperfect oxidation of one or more of the normal energy-producing food materials may be regarded as the basis of the so-called acidosis. It is distinguished by a typical group of symptoms, among which may be mentioned the elimination of the so-called acetone bodies - 3-hydroxybutyric acid, acetoacetic acid, and acetone - by the kidneys, a lessening of the titratable alkalinity of the blood, the hydrogen ion content remaining practically unchanged, a diminution of the carbon dioxide content of the alveolar air and a marked increase of the ammonia content of the urine. This group of symptoms is further found in certain other pathological and abnormal physiological states. For example, it is found that the feeding of mineral acids to herbivorous animals will produce a lessening of the titratable alkalinity of the blood, a diminution in the alveolar carbonic acid, and an increase in the urine of the fixed bases of the animal body, as sodium, potassium, calcium, and magnesium. The generally accepted explanation of this phenomenon, is that the acid on passing into the blood stream unites with the alkaline elements of the blood, and through the very delicate regulation of salt concentrations controlled by functional activity in the kidney, there is a definite loss of these alkaline elements, the normal function of which is to fix the carbon dioxide thrown off during the katabolic activity of the cells, and thus permit

it to be eliminated by the lungs. Similar experiments with the carnivora fail to exhibit a loss of the fixed alkalies, but do show a marked increase in the ammonia content of the urine. As the ash of the food of carnivora is acid in contradistinction to the alkaline ash of herbivorous nutrition, there is a partial protection normally established in the former organism. Further, the protein intake of the first far exceeds in quantity and utility that of the second, large quantities of ammonia being readily available through deaminization. Ammonia, then, may be regarded as the body's natural protection against increased acid formation. With man, a mixed feeder, we find the ammonia protection available, and thus in the relationship toward acid intoxications the eliminative features approach more nearly the picture of experimental acidosis with the carnivora.

Through disturbances in the organism we find the formation of varying amounts of the acetone bodies, and depending upon the origin, one or another symptom complex resulting from the same. For example, the lowering of the titratable alkalinity of the blood, with the appearance of the acetone bodies, the complementary lessening of the alveolar carbon dioxide, the appearance of the acetone bodies in the urine, and a marked increase in the urinary ammonia, form a condition which develops as a preliminary to diabetic coma. Further, we find

it to be eliminated by the lungs. Similar experiments with the carnivores fail to exhibit a loss of the fixed alkalies, but do show a marked increase in the ammonia content of the urine. As the ash of the food of carnivores is acid in contradistinction to the alkaline ash of herbivorous nutrition, there is a partial protection normally established in the former organism. Further, the protein intake of the first far exceeds in quantity and utility that of the second, large quantities of ammonia being readily available through decomposition. Ammonia, then, may be regarded as the body's natural protection against increased acid formation. With man, a mixed feeder, we find the ammonia protection available, and thus in the relationship toward acid intoxications the eliminative features approach more nearly the picture of experimental subjects with the carnivores.

Through disturbances in the organism we find the formation of varying amounts of the acetone bodies, and beginning upon the origin, one or another symptom complex resulting from the same. For example, the lowering of the titratable alkalinity of the blood, with the appearance of the acetone bodies, the complementary lessening of the alveolar carbon dioxide, the appearance of the acetone bodies in the urine, and a marked increase in the urinary ammonia, forms a condition which develops as a preliminary to diabetic coma. Further, we find

the comatose state in direct relationship with the amount of organic acid in the urine, coma appearing with increasing acid and disappearing with diminution of the acid content. The reduction of the alkalinity of the blood is an accompanying factor of the foregoing, and the fact frequently observed that the administration of alkalies tends toward improvement is in complete harmony with the facts already cited. On the other hand, in the disturbances of the organism produced by inanition, large quantities of B-hydroxybutyric acid will be eliminated by the urine without any evidence of coma. The source of this acid becomes a matter of primary interest. In diabetes the existing condition is one of a failure on the part of the organism to utilize available carbohydrates; in short, through one or another cause, a carbohydrate inanition exists. In the second instance, that of inanition, it is found that the glycogen reserve of the body is first drawn on for body maintenance and in a short time is materially exhausted. As before, the condition producing the acid intoxications rests upon a carbohydrate inanition. From this it seems safe to infer that the carbohydrates are literally anti-ketogenic in character, and that the acid formation cannot result from their imperfect oxidation. The other two possibilities, namely, the protein and fat of our nutrition, are potentially both sources of the acetone bodies. In regard to the former, we find the appearance of this acidosis associated with wasting diseases, in the course of

the comatose state in direct relationship with the amount of organic acid in the urine, coma appearing with increasing acid and disappearing with diminution of the acid content. The reduction of the alkalinity of the blood is an accompanying factor of the foregoing, and the fact frequently observed that the administration of alkalies tends toward improvement is in complete harmony with the facts already cited. On the other hand, in the disturbances of the organism produced by inanition, large quantities of B-hydroxybutyric acid will be eliminated by the urine without any evidence of coma. The source of this acid becomes a matter of primary interest. In diabetes the existing condition is one of a failure on the part of the organism to utilize available carbohydrates; in short, through one or another cause, a carbohydrate inanition exists. In the second instance, that of inanition, it is found that the glycogen reserve of the body is first drawn on for body maintenance and in a short time is materially exhausted. As before, the condition producing the acid intoxication rests upon a carbohydrate inanition. From this it seems safe to infer that the carbohydrates are literally anti-ketogenic in character, and that the acid formation cannot result from their imperfect oxidation. The other two possibilities, namely, the protein and fat of our nutrition, are potentially both sources of the acetone bodies. In regard to the former, we find the appearance of this acidosis associated with wasting diseases, in the course of

which large amounts of body protein are katabolized. On the other hand, more of the acetone bodies can be demonstrated in the urine in certain stages of the intoxication than can be accounted for by the protein destruction.

As regards the fats, experience teaches that not only would butyric acid produce the compounds in question, but that other higher fatty acids, if possessed of an even number of carbons, may also cause their formation. Further, laboratory experimentation has demonstrated that in the oxidation of fatty acids of large molecular weight, the molecules are progressively diminished by the loss of two carbons with each step in the decomposition. From this, it may be inferred that while both the protein and the fat are potentially sources of the acetone bodies, the fat seems to be the more probable source. The final solution of this phase of the problem, however, must be regarded as still in abeyance.

Differing from the group of acid intoxications, in which imperfect oxidation seems to be the causative factor, and in which the chief chemical features are the diminution of the alkalinity of the blood, and consequently of the alveolar carbonic acid, together with the elimination by the kidneys of the acetone bodies, and a complementary increase of the ammonia in the urine, is a group of pathological conditions, marked also by an acid intoxication, but in which functional disturbance of the liver seems to be the

which large amounts of body protein are metabolized. On the other hand, more of the acetone bodies can be formed in the urine in certain stages of the intoxication than can be accounted for by the protein destruction. As regards the fats, experience teaches that not only would butyric acid produce the symptoms in question, but that other higher fatty acids, if possessed of an even number of carbons, may also cause their formation. Further, laboratory experimentation has demonstrated that in the oxidation of fatty acids of large molecular weight, the molecules are progressively diminished by the loss of two carbons with each step in the decomposition. From this, it may be inferred that while both the protein and the fat are potentially sources of the acetone bodies, the fat seems to be the more probable source. The final solution of this phase of the problem, however, must be regarded as still in abeyance.

Turning from the group of acid intoxications, in which imperfect oxidation seems to be the causative factor, and in which the chief chemical features are the diminution of the alkalinity of the blood, and consequently of the alveolar carbonic acid, together with the elimination by the kidneys of the acetone bodies, and a complementary increase of the ammonia in the urine, is a group of pathological conditions, marked also by an acid intoxication, but in which functional disturbance of the liver seems to be the

its power of toleration of a carbohydrate deficiency.

The dominant cause. The intoxications by phosphorus and chloroform, the general toxemias of pregnancy, cyclic, pernicious vomiting, and acute yellow atrophy, are among the conditions falling in this second classification. Here, the acid is not of the group already mentioned, but is sarcolactic acid. This substance eliminated by the kidneys appears in the urine in liberal quantities, and associated with it is a marked increase, both in the ammonia content of the urine, and also that of the amino acids. That there is usually an accompanying diminution in the urea, points definitely to an impairment of the urea forming function of the liver, the ammonia and amino acids representing precursors of this final end product of general protein metabolism. In certain conditions, notably in the toxemias of pregnancy, there is, in addition to the lactic acid and the ammonia increase, an increase of the acetone bodies, so that the ammonia may be regarded as of two-fold origin. This is further borne out by the fact that the increase in ammonia is greater in the pernicious vomiting associated with pregnancy, than in eclampsia. That the latter is to be regarded more essentially as a toxemia, resting in part on disturbed liver function, while, in the former, imperfect oxidation also plays a significant role, is in perfect harmony with the facts presented above. Finally, the fact has been frequently observed that in what may be called normal pregnancy the organism seems to have lost, in part at least,

dominant cause. The intoxications by phosphorus and chloroform, the general toxemias of pregnancy, cyclic, pernicious vomiting, and acute yellow atrophy, are among the conditions falling in this second classification. Here, the acid is not of the group already mentioned, but is sarcosinic acid. This substance eliminated by the kidneys appears in the urine in liberal quantities, and associated with it is a marked increase, both in the ammonia content of the urine, and also that of the amino acids. That there is usually an accompanying diminution in the urea, points definitely to an impairment of the urea-forming function of the liver, the ammonia and amino acids representing precursors of this final end product of general protein metabolism. In certain conditions, notably in the toxemias of pregnancy, there is, in addition to the lactic acid and the ammonia increase, an increase of the acetone bodies, so that the ammonia may be regarded as of two-fold origin. This is further borne out by the fact that the increase in ammonia is greater in the pernicious vomiting associated with pregnancy, than in eclampsia. That the latter is to be regarded more essentially as a toxemia, resting in part on disturbed liver function, while, in the former, imperfect oxidation also plays a significant role, is in perfect harmony with the facts presented above. Finally, the fact has been frequently observed that in what may be called normal pregnancy the organism seems to have lost, in part at least,

its power of toleration of a carbohydrate deficiency. That the acidosis of the acetone bodies always causes a lessening of the alveolar carbon dioxide, and that the pernicious vomiting of pregnancy, in which this is a factor, is associated both with the non-convulsive toxemias and also with eclampsia, implies that the determination of the carbon dioxide elimination by the lungs should be of potential value, both from the diagnostic and the prognostic standpoint. Similarly, the determination of urinary ammonia, associated as it is with these conditions, is also to be regarded as of potential value in indicating the earlier stages of these types of acid intoxication. The chemical aspects of the determination of both ammonia and of carbon dioxide are varied and are affected by the imposition of certain limiting conditions. If the assumption may be made that the determination of one or the other of these is of sufficient value to justify its performance as a routine diagnostic procedure, then the adoption of a method, or of methods, which will give a sufficient degree of accuracy with a minimum of time consumption and complexity of apparatus and equipment, becomes a suitable matter for study. Since in the practical experience of every day, such determinations can be but minor incidents in the daily work of the busy medical practitioner, a study to be applicable to the needs of

its power of retention of a carbonyl group is relatively
That the action of the acetone bodies always remains
a lessening of the alveolar carbon dioxide, and that
the persistent vomiting of pregnancy, in which this is
a factor, is associated both with the non-convulsive
toxemia and also with eclampsia, implies that the deter-
mination of the carbon dioxide elimination by the lungs
should be of potential value, both from the diagnostic
and the prognostic standpoint. Similarly, the determi-
nation of urinary ammonia, associated as it is with these
conditions, is also to be regarded as of potential value
in indicating the earlier stages of these types of acid
intoxication. The chemical aspects of the determination
of both ammonia and of carbon dioxide are varied and are
affected by the imposition of certain limiting conditions.
If the assumption may be made that the determination of
one or the other of these is of sufficient value to justify
its performance as a routine diagnostic procedure, then
the adoption of a method, or of methods, which will give
a sufficient degree of accuracy with a minimum of time
consumption and complexity of apparatus and equipment,
becomes a suitable matter for study. Since in the prac-
tical experience of every day, such determinations can be
but minor incidents in the daily work of the busy medical
practitioner, a study to be applicable to the needs of

practice must include not only the consideration of sensitiveness, dependability, and accuracy, but also of simplicity and of time consumption.

The study then resolves itself into

- (1) a critical consideration of the various methods for the determination, respectively, of carbon dioxide in the alveolar air and of ammonia in the urine, leading to the selection of one or more methods for each which conform to the criteria established: and
- (2) by these methods, as adopted, to determine the relative dependability of the one or the other factor as an indication of approaching or incipient intoxication. To this could be suitably added continuous studies in those cases in which the toxemia failed of inhibition through treatment, to determine which could be regarded as the more suitable for following the progress of the case and giving the greater information for the prognosis. With these points in mind, the various available methods may now be subjected to critical analysis.

practice must include not only the consideration of sensitivity, dependability, and accuracy, but also of simplicity and of time consumption.

The study then resolves itself into

- (1) a critical consideration of the various methods for the determination, respectively, of carbon dioxide in the alveolar air and of ammonia in the urine, leading to the selection of one or more methods for each which conform to the criteria established; and
- (2) by these methods, as adopted, to determine the relative dependability of the one or the other factor as an indication of approaching or incipient intoxication. To this could be suitably added continuous studies in those cases in which the toxemia failed of inhibition through treatment, to determine which could be regarded as the more suitable for following the progress of the case and giving the greater information for the prognosis. With these points in mind, the various available methods may now be subjected to critical analysis.

AMMONIA DETERMINATIONS.

ALBIONIA DISTILLATIONS

AMMONIA DETERMINATIONS.

The methods of determination of urinary ammonia may be classified under two general headings: either it may be determined directly in the original solution by a wide variety of methods, or indirectly by removal in one way or another from the solution, the amount subsequently being estimated either by gravimetric, volumetric or colorimetric procedure.

The first course, determination in the original solution, offers many analytical possibilities; the ammonia nitrogen may be liberated as a gas by decomposition of its compounds and measured in suitable apparatus; other substances that might interfere with the colorimetric procedure may be removed and a method of direct Nesslerization and colorimetric determination employed; the ammonia may be precipitated by a suitable reagent in the form of a white cloud the amount being determined nephelometrically by comparison with known amounts similarly treated; or it may be determined indirectly by volumetric procedure according to the method best known as the Ronchese-Malfatti or formol titration.

The removal of the ammonia from the solution according to the second course, may likewise be accomplished in various ways. First, there is the aeration of the solution containing the ammonia compound the solution having been rendered strongly alkaline, (a) in the cold under atmospheric pressure, (b) the same, but under reduced pressure;

AMMONIA DETERMINATIONS.

The methods of determination of urinary ammonia

may be classified under two general headings: either it may be determined directly in the original solution by a wide variety of methods, or indirectly by removal in one way or another from the solution, the amount subsequently being estimated either by gravimetric, volumetric or colorimetric procedure.

The first course, determination in the original solution, offers many analytical possibilities; the ammonia nitrogen may be liberated as a gas by decomposition of its compounds and measured in suitable apparatus; other substances that might interfere with the colorimetric procedure may be removed and a method of direct measurement and colorimetric determination employed; the ammonia may be precipitated by a suitable reagent in the form of a white cloud the amount being determined nephelometrically by comparison with known amounts similarly treated; or it may be determined indirectly by volumetric procedure according to the method best known as the Nessler-Hafetz or formal titration.

The removal of the ammonia from the solution according to the second course, may likewise be accomplished in various ways. First, there is the separation of the solution containing the ammonia compound the solution having been rendered strongly alkaline. (a) in the cold under atmospheric pressure, (b) the same, but under reduced pressure;

second, there are the complementary methods where the ammonia is expelled from its compounds by heating either under normal or reduced pressure; and third, there is the precipitation and removal of ammonia by the use of a reagent with which it enters into chemical combination and from which it is subsequently liberated. In the aeration and distillation procedures the ammonia is in each case fixed by some acid solution placed in the receiver and the amount absorbed is determined gravimetrically by the formation of ammonium chloroplatinate or acid tartrate, volumetrically by titration of the excess of standard acid, or by the development of color with the Nessler-Winkler reagent and comparison with color produced under similar technique with known amounts of ammonia. The amount of the ammonia removed by precipitation may also be determined colorimetrically.

The following table classifies the various methods in a manner which may make clearer the subsequent analysis and criticism.

Determination of Ammonia.

I. Direct In Original Solution.

1. Decomposition of Ammonia.
 - a. Lematte
 - b. Florence
 - c. Bacon.
2. Direct Nesslerization.
 - a. Folin and Denis.
 - b. Sumner
3. Nephelometric
 - a. Graves
4. Direct Titration.
 - a. Ronchese-Malfatti

II. Indirect After Removal from Solution.

1. Vacuum Distillation.

- a. Boussingault
- b. Würster
- c. Nencki-Zaleski and Soldern
- d. Steyer
- e. Schaffer-Titration
- f. Krüger Reich Schittenhelm

2. Distillation.

- a. Schlösing-Titration
- b. deGraaff-Titration
- c. Bonnema-Titration
- d. Winkler-Titration

3. Aeration.

- a. Folin-Titration
- b. Kober-Titration
- c. Davis
- d. Folin and MacCallum Titration or Colorimeter

4. Precipitation.

- a. Folin and Bell-Colorimeter.

The determination of ammonia by decomposition according to the methods outlined by Lematte (1), Florence (2) and Bacon (3), is at best indirect. The methods involve the determination of urea plus ammonia and the determination of urea alone, the ammonia value being determined by difference. Lematte precipitates the ammonia salts with phosphotungstic acid and magnesium chloride and determines the urea in the filtrate by the hypobromite method. Florence removes Uric acid and Creatinine with basic lead acetate and decomposes the ammonia salts by heating with the same reagent and determines the urea in resulting solution by the hypobromite method. Bacon uses Millon's reagent which yields Nitrogen and carbon dioxide in equal volume but does not decompose ammonia or its salts. This reaction like the hypo-

- | | | |
|----------------|---------------------------|--------|
| 1. L. Lematte | Comp. Rend. Soc. de Biol. | (1913) |
| 2. A. Florence | " " " " | (1910) |
| 3. R.F. Bacon | Phil. Jr. of Science. | (1909) |

II. Indirect After Removal from Solution.

1. Vacuum Distillation.

- a. Bousignault
- b. Winter
- c. Nencki-Salski and Solberg
- d. Steyer
- e. Scheller-Titration
- f. Krüger, Reichen, Schreinemakers

2. Distillation.

- a. Scheller-Titration
- b. Scheller-Titration
- c. Scheller-Titration
- d. Scheller-Titration

3. Titration.

- a. Folin-Titration
- b. Folin-Titration
- c. Davis
- d. Folin and MacCallum Titration or Colorimeter

4. Precipitation.

- a. Folin and Balli-Colorimeter.

The determination of ammonia by decomposition

according to the methods outlined by Lematta (1), Florence (2)

and Bacon (3), is at best indirect. The methods involve

the determination of urea plus ammonia and the determination

of urea alone, the ammonia value being determined by differ-

ence. Lematta precipitates the ammonia salts with phospho-

tungstic acid and magnesium chloride and determines the urea

in the filtrate by the hypobromite method. Florence removes

Uric acid and creatinine with basic lead acetate and decom-

poses the ammonia salts by heating with the same reagent and

determines the urea in resulting solution by the hypobro-

mite method. Bacon uses Millon's reagent which yields

nitrogen and carbon dioxide in equal volume but does not de-

compose ammonia or its salts. This reaction like the hypo-

1. I. Lematta	Comp. Rend. Soc. de Biol.	(1913)
2. A. Florence	" " " "	(1910)
3. F. Bacon	Brit. J. of Hygiene.	(1909)

bromite of the other two is not quantitative, approximate results only being obtained. According to all three procedures the urea plus ammonia value is determined by means of hypobromite methods. The basis of the various hypobromite methods depends upon the fact that urea and ammonia-containing compounds are decomposed by a solution of sodium hypobromite with the formation of carbon dioxide and gaseous nitrogen. The carbon dioxide is retained by the excess alkali in the solution and the nitrogen gas is collected and measured. The method when carried out with the greatest of care, avoiding all possible mechanical errors due to effects of temperature, lack of alkali or reagent, loss of gas, and incorrect graduation of the apparatus, is at best but a clinical method for the estimation of urea and ammonia nitrogen, and is in no sense a strictly quantitative determination. Robinson and Muller have made comparative studies of the method using the Doremus-Hinds Ureometer and comparing it with the Folin-Pettibone Phosphoric acid method. Their conclusions are that the method gives inconsistent and completely unreliable results when performed in the usual way as described in textbooks, a constant reading not being obtained for from five to twenty-four hours after application of the reagent. They also find that uncorrected instruments give readings which vary from ninety-seven to one hundred and five percent of the correct value. In the presence of glucose in the urines there is an apparent catalysis and the readings are much higher than those of normal urines.

proximate of the other two is not quantitative, approximate results only being obtained. According to all three procedures the true value is determined by means of hypochlorite method. The basis of the various hypochlorite methods depends upon the fact that urea and ammonia-containing compounds are decomposed by a solution of sodium hypochlorite with the formation of carbon dioxide and gaseous nitrogen. The carbon dioxide is retained by the excess alkali in the solution and the nitrogen gas is collected and measured. The method when carried out with the greatest of care, avoiding all possible mechanical errors due to effects of temperature, lack of alkali or reagent, loss of gas, and incorrect graduation of the apparatus, is at best but a clinical method for the estimation of urea and ammonia nitrogen, and is no more a strictly quantitative determination. Johnson and Miller have made comparative studies of the method using the Gormann-Hinds Ureometer and comparing it with the Folin-Cutcliffe Phosphoric acid method. Their conclusions are that the method gives inconsistent and completely unreliable results when performed in the usual way as described in textbooks, a constant reading not being obtained for five to twenty-four hours after application of the reagent. They also find that micro-tested instruments give readings which vary from ninety-seven to one hundred and five percent of the correct value. In the presence of glucose in the urine there is an apparent catalysis and the readings are much higher than those of normal urines.

Since the ammonia content is determined by difference, the errors incident to the two reactions exercise the maximum influence upon its absolute magnitude. Further, as under the most favorable conditions the ammonia nitrogen is but a relatively small fraction of the total amount determined, the error not infrequently assumes wholly vitiating proportions.

The methods for direct Nesslerization are two; the Merck's blood charcoal method suggested by Folin and Denis(1), and the copper hydroxide method described by Sumner (2). The first method depends upon the use of blood charcoal to remove by adsorption the greater part of the creatinine, Uric acid and pigmentary bodies, all of which would interfere with the colorimetric determination. The procedure is in brief as follows: To 10cc of Urine in a flask is added 1cc of 25% m-phosphoric acid, 9cc distilled water and 2gm. of blood charcoal. Shake well for at least one minute and filter through a dry paper. Transfer 1 to 5cc of the filtrate to a 100cc volumetric flask, add distilled water to 70cc Nesslerize by the addition of 15cc Nessler's solution, make up to volume with distilled water and compare with a solution obtained by Nesslerizing 1 mg. of ammonia nitrogen in another 100cc flask. The method is rapid and reasonably accurate. As used by its authors it gives results which

- | | |
|--------------------------|---------------------------|
| 1. O. Folin and W. Denis | Jr. of Biol. Chem. (1916) |
| 2. J.B.Sumner | " " " " (1918) |

are comparable with those obtained by the Folin-MaCallum macro-aeration method. The accuracy of the method, however, is dependent upon the efficiency of the reagent which is a highly variable quantity. Moreover, none but Merck's purified blood charcoal has been found to have the necessary properties, and that reagent is not made in this country and cannot be obtained.

In the second method cupric hydroxide is made use of to precipitate the creatinine. The technique of the method is as follows: Pipette into a large test tube 10cc saturated copper sulphate solution, 15cc of Urine and 10cc of sodium hydroxide (Normality 2.03). Stopper. Shake. Filter, covering the funnel to prevent evaporation. Nesslerize a portion of the filtrate and compare in the colorimeter against a standard containing one milligram of nitrogen. The author's tabulated results show the method to be comparable with the aeration procedure. The sodium hydroxide solution should be free both from iron and carbonate and the copper sulphate must be free from ammonia. Too, in either an acid or an alkaline medium the precipitation of both creatinine and copper is interfered with. At the neutral point 90% of the Creatinine will be carried down.

The Nephelometric determination as suggested by Graves (1) as a substitute for the colorimetric determination with the Nessler-Winkler reagent is a relatively unimportant method. It requires the use of apparatus that is not commonly found in the laboratory and offers no ad-

are comparable with those obtained by the Volin-MacAllum
macro-estimation method. The accuracy of the method, how-
ever, is dependent upon the efficiency of the reagent
which is a highly variable quantity. Moreover, none but
Morck's purified blood charcoal has been found to have the
necessary properties, and that reagent is not made in this
country and cannot be obtained.

In the second method cupric hydroxide is made use
of to precipitate the creatinine. The technique of the
method is as follows: Pipette into a large test tube 10cc
saturated copper sulphate solution, 15cc of urine and 10cc
of sodium hydroxide (Normality 2.05). Stopper. Shake.
Filter, covering the funnel to prevent evaporation. Measure
a portion of the filtrate and compare in the colorimeter
against a standard containing one milligram of nitrogen. The
author's tabulated results show the method to be comparable
with the reaction procedure. The sodium hydroxide solution
should be free from iron and carbonate and the copper
sulphate must be free from ammonia. Too, in either an acid
or an alkaline medium the precipitation of both creatinine
and copper is interfered with. At the neutral point 90%
of the creatinine will be carried down.

The Nephelometric determination as suggested by
Graves (1) as a substitute for the colorimetric determin-
ation with the Nessler-Linkler reagent is a relatively
unimportant method. It requires the use of apparatus that
is not commonly found in the laboratory and offers no ad-

advantages over the colorimetric method. He suggests the precipitation of ammonia from the solution by means of a reagent prepared from 50cc HgCl_2 , 15 gm NaCl , 35cc saturated Li_2CO_3 solution and 65cc of water, and its subsequent comparison with similarly precipitated known amounts of ammonia.

The last of these direct determinations in the original solution is the Ronchese-Malfatti (1) formalin titration which offers excellent possibilities, and while the general consensus of opinion places it among the relatively inaccurate methods, a careful analysis of the probable sources of error shows that several of these may readily be corrected. This permits of an accuracy with which the method has never been credited. The method may be briefly described as follows:

A standard sample of the urine, usually 25 cubic centimeters, is rendered exactly neutral to phenolphthalein by the addition of tenth-normal sodium hydroxide. Diluting the urine with distilled water sharpens the end point, and the addition of solid potassium oxalate, as suggested by Polin, eliminates the disturbing effect of the calcium salts. To the neutral urine is next added a definite amount of neutral commercial formaldehyde solution. It is imperative that the neutrality of this latter be insured, as the commercial product always contains appreciable amounts of formic acid, and the auto-oxidative process goes

1. A. Ronchese Comp. Rend. Soc. de Biol. (1907)
2. H. Malfatti Zeit. f. Anal. Chem. (1908)

on whenever the solution is exposed to the air. To the neutral solution thus obtained, tenth-normal caustic soda is run in until the appearance of the pink color due to the phenolphthalein shows that the neutral point has been attained. The method depends upon the fact that the caustic soda liberates ammonia from its several compounds, and that the latter, in the presence of formaldehyde, reacts quantitatively to form hexamethylenetetramine, a compound which is neutral toward phenolphthalein. The volume of caustic soda determines at once the amount of ammonia present.

One source of error in this method rests upon the fact that amino groups show the same reaction as do ammonia compounds. Under ordinary conditions, the amino acid content of the urine is so small as to make unlikely their forming a disturbing factor in the method. In, however, conditions of functional impairment of the liver, where the deaminizing power of the organ is impaired, the amino acid content of the urine increases to appreciable proportions, and thus forms a serious source of error in the direct ammonia determination. In consideration of the fact that the ammonia increase is, in part, due to the same cause as that which increased the amino acids: namely, an impairment of the ureagenetic function, the significance of the one compound is essentially that of the other. The significance of a marked increase in either or both would be the same. For the purposes of this study, then, this amino acid error becomes relatively insignificant.

Another source of error, which seems to have been ignored in the literature, but which offers an absolute and variable element of uncertainty, lies in the carbonate content of the caustic soda solution, and of the urine. Since phenolphthalein, a substance easily affected by carbonates has been selected as the indicator, it is obvious that unless this factor can be eliminated, the possible error of the method is increased. The alkali should be prepared in the carbonate-free state, and then maintained free from carbon dioxide contamination. The assumption that high-grade, alcohol-purified sodium hydroxide is free from carbonate contamination may be demonstrated to be erroneous. The best grades of caustic soda on the market will show from two to five percent of carbonate. If to this is added the carbon dioxide contained in the average laboratory supply of distilled water, the amount is further increased. The procedure followed in this laboratory is as follows:

The caustic soda solution is prepared at slightly greater strength than in the final concentration desired. The carbonate content is determined by parallel titrations at zero degree, using respectively phenolphthalein and methyl orange as indicators. Then slightly more barium hydroxide than the calculated amount necessary to precipitate the carbonate is added. After the carbonate precipitate has settled, the clear liquid is decanted in a carbon dioxide-free atmosphere, and the barium content determined

Another source of error, which seems to have been ignored in the literature, but which offers an absolute and variable element of uncertainty, lies in the carbonate content

of the caustic soda solution, and of the urine. Since phenolphthalein, a substance easily affected by carbonates, has been selected as the indicator, it is obvious that unless this factor can be eliminated, the possible error of

the method is increased. The alkali should be prepared in the carbonate-free state, and then maintained free from carbon dioxide contamination. The assumption that high-

grade, alcohol-purified sodium hydroxide is free from carbonate contamination may be demonstrated to be erroneous. The best grades of caustic soda on the market will show from two to five percent of carbonate. If to this is added the carbon dioxide contained in the average laboratory supply of distilled water, the amount is further increased. The procedure followed in this laboratory is as follows:

The caustic soda solution is prepared at slightly greater strength than in the final concentration desired. The carbonate content is determined by parallel titrations at zero degree, using respectively phenolphthalein and methyl orange as indicators. Then slightly more barium hydroxide than the calculated amount necessary to precipitate the carbonate is added. After the carbonate precipitate has settled, the clear liquid is decanted in a carbon dioxide-free atmosphere, and the barium content determined

gravimetrically as barium sulphate. The calculated amount of sulphuric acid, or better sodium sulphate, is added, and after the subsidence of the barium sulphate precipitate, the carbonate and barium-free alkali solution is transferred in a carbon dioxide-free atmosphere to a paraffin-lined stock bottle. The use of absorbing towers filled with soda lime for the purification of all air entering the stock bottle maintains the liquid free from carbonate contamination.

The second source of carbonate contamination: namely, the carbon dioxide in the urine, can be readily eliminated by the method of procedure. The initial neutralization of the urine is affected by the carbonate content, thereby introducing an error into the determination of the titratable acidity, which constitutes this initial step. This fact has not been touched upon in the literature, so far as the author is aware, and experiments made in this laboratory show that errors of appreciable magnitude may arise in this connection. The urine being neutral toward phenolphthalein after the completion of this step, the second titration merely involves a re-establishment of the same P_h value, and if the alkali be free from carbonate the amount used is a measure of the ammonia content.

When we come to the indirect determination of ammonia the first problem is the liberation of ammonia from its compounds in the urine without causing the decomposition of other nitrogen containing substances present

gravimetrically as barium sulphate. The calculated amount of sulphuric acid, or better sodium sulphate, is added, and after the subsidence of the barium sulphate precipitate, the carbonate and barium-free alkaline solution is transferred in a carbon dioxide-free atmosphere to a carefully-lined stock bottle. The use of absorbing towers filled with soda lime for the purification of all air entering the stock bottle maintains the liquid free from carbonate contamination.

The second source of carbonate contamination; namely, the carbon dioxide in the urine, can be readily

eliminated by the method of procedure. The initial neutralization of the urine is effected by the carbonate content, thereby introducing an error into the determination of the titratable acidity, which constitutes this initial step. This fact has not been pointed out in the literature, so far as the author is aware, and experiments made in this laboratory show that errors of appreciable magnitude may arise in this connection. The urine being neutral toward phenolphthalein after the completion of this step, the second titration merely involves a re-establishment of the same pH value, and if the alkali be free from carbonate the amount used is a measure of the ammonia content.

When we come to the indirect determination of ammonia the first problem is the liberation of ammonia from its compounds in the urine without causing the decomposition of other nitrogen containing substances present

there. Various alkalis have been tried. Milk of lime and magnesia were used in the early methods of Schlösing (1) Boussingault (2) and Würster (3). The lime gives off the ammonia rapidly but causes the decomposition of other substances resulting in very high ammonia values. The magnesia on the other hand gives off the ammonia very slowly and the removal is incomplete, due to the formation of ammonium-magnesium phosphate crystals from which the ammonia cannot be liberated by magnesia. Boussingault also made use of sodium carbonate the reagent which has since been suggested by Folin (4) and Schaffer (5). This reagent drives off the ammonia more rapidly than does the magnesia and does not cause decomposition as the lime does. It has, however, been shown by Steele (6) and by S. R. Benedict (7) that in the presence of magnesium in the urine, the addition of carbonate and the process of aeration will cause the formation of ammonium-magnesium phosphate crystals. Steele and Gies (8) suggest the use of sodium hydroxide for the purpose of liberating the ammonia and the addition of sodium chloride to force back the dissociation of the hydroxide and render its action less drastic. The experimental evidence offered by both Steele and Benedict demonstrates

1. Schlösing Annales de Chem. et de Phys. (1851)
2. Boussingault " " " " " (1850)
3. C. Würster Central f. Phys. (1887)
4. O. Folin Zeit für Phys. Chem. (1902)
5. P. Schaffer Am. Jr. of Phys. (1903)
6. Steele, Proc. Soc. Exp. Biol. (1909)
7. S. R. Benedict and Emil Osterberg, Biochem. Bull. (1913)
8. Steele and Gies, Jr. Biol. Chem. (1909)

there. Various alkalis have been tried. Milk of lime and magnesia were used in the early methods of Schilling (1) and Boussingault (2) and Whistler (3). The lime gives off the ammonia rapidly but causes the decomposition of other substances resulting in very high ammonia values. The magnesia on the other hand gives off the ammonia very slowly and the removal is incomplete, due to the formation of ammonium-magnesium phosphate crystals from which the ammonia cannot be liberated by magnesia. Boussingault also made use of sodium carbonate the reagent which has since been suggested by Folin (4) and Schaffer (5). This reagent drives off the ammonia more rapidly than does the magnesia and does not cause decomposition as the lime does. It has, however, been shown by Steele (6) and by A. R. Benedict (7) that in the presence of magnesium in the urine, the addition of carbonate and the process of aeration will cause the formation of ammonium-magnesium phosphate crystals. Steele and Gies (8) suggest the use of sodium hydroxide for the purpose of liberating the ammonia and the addition of sodium chloride to force back the dissociation of the hydroxide and render its action less drastic. The experimental evidence offered by both Steele and Benedict demonstrates

- | | |
|--------------------------------------|-------------------------------------|
| 1. Schilling | Annales de Chem. et de Phys. (1851) |
| 2. Boussingault | " " " " (1850) |
| 3. C. Whistler | Central J. Phys. (1867) |
| 4. O. Folin | Zeit für Phys. Chem. (1908) |
| 5. F. Schaffer | Am. J. of Phys. (1908) |
| 6. Steele | Proc. Soc. Biol. (1909) |
| 7. A. R. Benedict and Emil Osterberg | Biochem. Bull. (1912) |
| 8. Steele and Gies | Jr. Biol. Chem. (1909) |

The matter will be discussed further in connection with that this reagent will liberate ammonia quantitatively from the triple phosphate and that other substances are not decomposed by the treatment so as to yield ammonia. Folin (1) preferring sodium carbonate to sodium hydroxide modified his method by adding acid to dissolve the crystals and from seven to eleven grams of potassium oxalate to the sodium carbonate. This modification Benedict finds is not so satisfactory in the original Folin method as is the modification by Steele and Gies, tho in the macro-aeration method where the amount of oxalate necessary is decreased, the method is successful. Potassium hydroxide cannot be used to liberate the ammonia for potassium sulphate if formed may carry down ammonia by occlusion as a double salt.

The ammonia driven out of solution in all distillation and aeration procedures is fixed by some acid solution placed in the receiver. The amount of ammonia is then determined by titration of the excess acid, by the development of color with the Nessler-Winkler reagent or as suggested by Foxwell (2) with phenol and sodium chlorate, or gravimetrically by the preparation of an insoluble ammonium salt as chloroplatinate or acid tartrate. The acid used in the collection is usually sulphuric; hydrochloric, however, is used by Folin and McCallum (3) in their aeration procedure and Winkler modifies the Kjeldahl procedure by the use of Boric acid. The relative merits of titration and colorimetric determinations are still a matter of controversy among the advocates of one or the other procedure.

1. O. Folin, Jr. Biol. Chem. (1910)
2. Foxwell, Gas World, (1916)
3. O. Folin and MacCallum, Jr. Biol. Chem. (1912)

that this reagent will liberate ammonia quantitatively from the triple phosphate and that other substances are not decomposed by the treatment so as to yield ammonia. Folin (1) preferring sodium carbonate to sodium hydroxide modified his method by adding acid to dissolve the crystals and from seven to eleven grams of potassium oxalate to the sodium carbonate. This modification Benedict finds is not so satisfactory in the original Folin method as is the modification by Steele and Cies. This is the macro-estimation method where the amount of oxalate necessary is decreased, the method is successful. Potassium hydroxide cannot be used to liberate the ammonia for potassium sulphate is formed very heavy brown ammonia

by oxidation as a foshie salt.

The ammonia driven out of solution in all distillation and reaction procedures is fixed by some acid solution placed in the receiver. The amount of ammonia is then determined by titration of the excess acid, by the development of color with the Nessler-Winkler reagent or an reagent by Foxwell (2) with phenol and sodium chloride, or gravimetrically by the preparation of an insoluble ammonium salt as calcium carbonate or acid tartrate. The acid used in the collection is usually sulphuric; hydrochloric, however, is used by Folin and McCaffrey (3) in their reaction procedure and Winkler modifies the Kjeldahl procedure by the use of boric acid. The relative merits of titration and colorimetric determinations are still a matter of controversy among the advocates of one or the other procedure.

The matter will be discussed further in connection with the methods involving their use.

The colorimetric method as suggested by Foxwell, has not, so far as the writer knows, been applied to urinary determinations. He treats the ammonia solution with phenol and sodium chlorate and compares it in a small colorimeter, such as is used in the Marriott carbon dioxide determination, with previously prepared standard solutions containing varying amounts of ammonia similarly treated with phenol and sodium chlorate. The method is potentially useful as a clinical procedure in that it obviates the necessity of the purchase of an expensive colorimeter which of course is necessary for the Nessler-Winkler comparisons. The length of time that the standards prepared for the Foxwell comparison would retain their integrity, is a factor that would enter into the evaluation of the method.

The gravimetric determination is susceptible of certain unavoidable errors which when linked to the elaborate and time consuming technique of the method makes it obviously unsuited for the problem under consideration.

Vacuum distillation is, so far as the writer can find, the earliest method used for the removal of ammonia from the urine. In 1850 Boussingault published a method for determining ammonia by distilling to dryness with milk of lime in a vacuum giving 40 to 50 degrees Centigrade. The operation took about an hour. This, tho undoubtedly the original application of vacuum distillation to the determination of ammonia, appears to have received little attention. Würster in 1887 proposed a similar method for

The matter will be discussed further in connection with

the methods involving their use.

The colorimetric method as suggested by Foxwell,

has not, so far as the writer knows, been applied to urinary

determinations. He treats the ammonia solution with phenol

and sodium chlorate and compares it in a small colorimeter,

such as is used in the Harriott carbon dioxide determination,

with previously prepared standard solutions containing vary-

ing amounts of ammonia similarly treated with phenol and

sodium chlorate. The method is potentially useful as a

clinical procedure in that it obviates the necessity of the

purchase of an expensive colorimeter which of course is nec-

essary for the Nessler-Winkler comparisons. The length of

time that the standards prepared for the Foxwell comparison

would retain their intensity, is a factor that would enter

into the evaluation of the method.

The gravimetric determination is susceptible of

certain unavoidable errors which when linked to the elab-

orate and time consuming technique of the method makes it

obviously unsuited for the problem under consideration.

Vacuum distillation is, so far as the writer can

find, the earliest method used for the removal of ammonia

from the urine. In 1850 Boissacq published a method

for determining ammonia by distilling to dryness with milk

of lime in a vacuum giving 40 to 50 degrees Centigrade.

The operation took about an hour. This, the undoubtedly

the original application of vacuum distillation to the

determination of ammonia, appears to have received little

attention. Whistler in 1887 proposed a similar method for

distillation and apparently was not aware of Boussingault's work. Würster (1) distilled at fifteen millimeters pressure at 50 degrees Centigrade using five, ten or twenty cc of Urine with baryta water, magnesia or lime, and claimed the operation to be completed after five, ten or fifteen minutes depending upon the amount of urine used. The method is inaccurate due to the small quantities of urine used and to the incompleteness of the liberation of ammonia under the conditions of the determination. The results while acceptable in the last century, are not to be compared with the accuracy of present day methods. The apparatus used was bulky and inconvenient. Modifications of the method were made by Nencki (2), Zaleski (2) and Soldern (3) and by Steyer (4), modifications all of which sought to construct an apparatus by various arrangements of flasks which would obviate the loss of the determination by the foam getting into the alkali. The modifications increased the accuracy also by increasing the quantity of urine and continuing the distillation over a longer period of time.

In 1903 Schaffer (5) made a modification of the original Boussingault method: In order to shorten the period of distillation and so lessen the error due to decomposition without using too small amounts of the urine Schaffer added methyl alcohol to the urine thus lowering

1. C. Würster, Central. f. Phys. (1887)
2. M. Nencki and J. Zaleski, Archiv. f. Exper. Path. (1895)
3. Soldern, Zeit f. Biol. (1899)
4. Steyer, Beitr. zur Chem. Phys. u. Path. (1902)
5. Schaffer, Am. Jr. of Phys. (1903)

distillation and apparently was not aware of Boussingault's work. Whistler (1) distilled at fifteen millimeters pressure at 50 degrees Centigrade using five, ten or twenty cc of urine with baryta water, ammonia or lime, and claimed the operation to be completed after five, ten or fifteen minutes depending upon the amount of urine used. The method is inaccurate due to the small quantities of urine used and to the inconstancy of the liberation of ammonia under the conditions of the determination. The results while acceptable in the last century, are not to be compared with the accuracy of present day methods. The apparatus used was bulky and inconvenient. Modifications of the method were made by Mensch (2), Salanski (3) and Solheim (3) and by Steyer (4). Modifications all of which sought to construct an apparatus by various arrangements of flasks which would obviate the loss of the determination by the foam getting into the alkali. The modifications increased the accuracy also by increasing the quantity of urine and containing the distillation over a longer period of time. In 1905 Schaeffer (5) made a modification of the original Boussingault method. In order to shorten the period of distillation and to lessen the error due to decomposition without using too small amounts of the urine Schaeffer added methyl alcohol to the urine thus lowering

1. C. Whistler, Central. E. Phys. (1867)
2. M. Mensch and J. Salanski, Archiv. E. Exper. Path. (1895)
3. Solheim, Zeit. E. Biol. (1899)
4. Steyer, Beitr. zur Chem. Phys. u. Path. (1902)
5. Schaeffer, An. Ch. de Phys. (1905)

the boiling point so that a more rapid ebullition would take place. He reduced the time to fifteen minutes, using sodium carbonate to liberate the ammonia. In the same year, Kruger, Reich and Schittenhelm (1) published a similar method using sodium carbonate and sodium chloride and added ethyl alcohol to prevent foaming.

The method of Schaffer may be considered as typical of this type of determination. It is in brief as follows: To 50cc of urine in a flask, add an excess of sodium chloride and about 50cc of methyl alcohol. This is connected with two absorption flasks in series, the first containing 50cc, the second 10cc of standard tenth normal acid. The second bottle is in turn connected through a safety flask with a vacuum pump. The urine container is placed in a water bath at 50° and one gram of dry sodium carbonate added; suction is then started and the boiling continued for fifteen minutes. The acid in the two absorption flasks is then titrated, using Alizarin as the indicator.

A distillation method without the use of a vacuum was described by Schlösing (2) in 1851 and was used to determine the ammonia in tobacco. In 1898 Neubauer (3) in his "Anteilung für Harn Analyse" applied the method to urine. As described by Neubauer the method was as follows: To 10cc of urine was added magnesia or milk of lime and this mixture was placed in a flat dish ten or twelve centimeters wide; over

1. Kruger, Reich and Schittenhelm, Jr. f. Phys. Chem. (1903)
2. Schlösing, Annal. de. Chem. et de Phys. (1851)
3. Neubauer, Jr. f. Prak. Chem. (1855)

The boiling point is that of pure rapid evaporation would
take place. He reduced the time to fifteen minutes.
using sodium carbonate to liberate the ammonia. In the
same year, Wray, Eichen and Schilling (1) published
a similar method using sodium carbonate and sodium chloride
and which they claimed to prevent foaming.

The method of Schilling may be considered as
typical of this type of determination. It is in brief
as follows: To 50cc of urine in a flask, add an excess of
sodium chloride and about 50cc of methyl alcohol. This
is connected with two absorption flasks in series, the
first containing 50cc, the second 100cc of standard ferrous
normal acid. The second bottle is then connected
through a safety flask with a vacuum pump. The urine con-
tainer is placed in a water bath at 50° and one gram of
dry sodium carbonate added; ammonia is then evolved and
the boiling continued for fifteen minutes. The acid in
the two absorption flasks is then titrated, using potassium
as the indicator.

A distillation method without the use of a vacuum
was described by Schilling (2) in 1931 and was used to de-
termine the ammonia in tobacco. In 1933 Kohnen (3) in
his "Analytische Ursubstanzanalyse" applied the method to urine.
It is described by Kohnen as follows: To 100cc
of urine was added ammonia or milk of lime and this mixture
was placed in a 250cc flask or twelve centimeter wide; over

1. Wray, Eichen and Schilling, J. E. Chem. Soc. (1933)
2. Schilling, Annal. der Chem. 464 (1931)
3. Kohnen, J. E. Prakt. Chem. (1933)

this dish and supported by a triangle was another dish and supported by a triangle was another dish containing the acid; all was closed by a tightly fitted bell jar and allowed to stand forty-eight hours. The authors used standard acids and alkali of such strength as to make it impossible to measure the small quantities of ammonia given off on standing longer than forty-eight hours and so considered their determinations complete. The method is susceptible of many errors and is decidedly unreliable.

Heat distillations using modifications of the Kjeldahl apparatus have been suggested by other chemists. De Graaff and Bonnema (1) distil into N/10 acid using sodium carbonate and calcium oxide respectively to liberate the ammonia. Ethyl alcohol is added to prevent foaming. Winkler uses Boric acid for the absorption of the ammonia and titrates with standard hydrochloric or sulphuric acid using Congo red or Methyl orange as indicators. The method is of necessity not an exact one. Boric acid is too weak an acid to completely absorb the ammonia and the end point in a titration with either Congo red or Methyl orange is not sharp.

Folin in 1903 worked out the first of the aeration methods as a substitute for distillation. The procedure in brief is as follows: Place 25cc of urine in an aerometer cylinder, add about 1 gm. of dry sodium carbonate and introduce some crude petroleum to prevent foaming. Connect with the suction pump through a cal-

- | | | |
|--------------|-----------------|--------|
| 1. DeGraaff, | Pharm. Weekblad | (1916) |
| Bonnema, | Chem. Ztg. | (1915) |

this dish and supported by a triangle was another dish and supported by a triangle was another dish containing the acid; all was closed by a tightly fitted bell jar and allowed to stand forty-eight hours. The authors used standard acids and alkali of such strength as to make it impossible to measure the small quantities of ammonia given off on standing longer than forty-eight hours and so considered their determinations complete. The method is susceptible of many errors and is decidedly unreliable.

Heat distillations using modifications of the Kjeldahl apparatus have been suggested by other chemists. De Graaf and Bonnera (1) distill into $N/10$ acid using sodium carbonate and calcium oxide respectively to liberate the ammonia. Ethyl alcohol is added to prevent foaming. Finkler uses boric acid for the absorption of the ammonia and titrates with standard hydrochloric or sulphuric acid using Congo red or Methyl orange as indicators. The method is of necessity not an exact one. Boric acid is too weak an acid to completely absorb the ammonia and the end point in a titration with either Congo red or Methyl orange is not sharp.

Polin in 1903 worked out the first of the aeration methods as a substitute for distillation. The procedure in brief is as follows: Place 25cc of urine in an aerometer cylinder, add about 1 gm. of dry sodium carbonate and introduce some crude petroleum to prevent foaming. Connect with the suction pump through a cal-

cium chloride tube filled with cotton and an absorbing flask containing 20cc of N/10 sulphuric acid, ammonia free water and a few drops of alizarin. With an ordinary filter pump and good water pressure the last trace of ammonia should be removed in about an hour and a half. The number of cubic centimeters of the N/10 acid neutralized by the ammonia of the urine may be determined by direct titration with N/10 sodium hydroxide. The modification in the use of a reagent suggested by Steele has already been discussed. Folin recommends the aeration procedure as a substitute for Kjeldahl nitrogen determination. Whether or not it is an acceptable substitute has been the subject of some controversy. If there is an adequate volume of air, a small volume of liquid and an excess of alkali, the method is susceptible of a high degree of accuracy. Kober (1) in 1909 and Davis (2) in 1909 describe aeration methods similar in technique to that of Folin.

The Folin-MaCallum (3) micro-aeration method which is a modification of the original Folin is a combination of the aeration procedure for ammonia with its colorimetric determination by means of the Nessler-Winkler reagent. Because small quantities of urine are used, thus shortening the time necessary for aeration, the method is considered by its authors available for clinical determinations. From one to five cc of the urine are placed in a test-tube, a few drops of a solution of potassium carbonate and potassium

- | | |
|------------------------|-----------------------------|
| 1. P.A.Kober, | Jr. Amer. Chem. Soc. (1913) |
| 2. R.O.Davis, | " " " " (1909) |
| 3. Folin and MaCallum, | Jr. Biol. Chem. (1912) |

clear solution was filled with cotton wool and an absorbing
flask containing 200 cc of 10% sulphuric acid, ammonia
free water and a few drops of indicator. With an orifice
very little pump and good water pressure the first traces
of ammonia should be removed in about an hour and a half.
The number of cubic centimeters of the 10% acid solution
required by the ammonia of the urine may be determined
by direct titration with 10% sodium hydroxide. The
addition in the use of a reagent suggested by Stalla
has already been discussed. This reagent is the same
as the procedure as a substitute for Kjeldahl nitrogen
determination. Whether or not it is an acceptable sub-
stitute has been the subject of some controversy. It there
is an adequate volume of air, a small volume of liquid and
an excess of alkali, the method is susceptible of a high
degree of accuracy. Kober (1) in 1903 and Davis (2) in
1903 describe reaction methods similar in technique to that
of Folin.

The Folin-MacCallum (3) micro-reaction method
which is a modification of the original Folin is a combina-
tion of the reaction procedure for ammonia with its color-
imetric determination by means of the Fessler-Linkler re-
agent. Because small quantities of urine are used, this short-
ening the time necessary for reaction, the method is consid-
ered by the authors available for clinical determinations.
From one to five cc of the urine are placed in a test-tube,
few drops of a solution of potassium carbonate and potassium

- | | |
|-------------------------|-----------------------------|
| 1. F.A. Kober, | Tr. Amer. Chem. Soc. (1913) |
| 2. R.D. Davis, | " " (1903) |
| 3. Folin and MacCallum, | Tr. Biol. Chem. (1913) |

oxalate are added, and a strong air current is passed through the mixture until the ammonia has all been removed. The ammonia is collected in ammonia-free water and acid, is Nesslerized, and the amounts determined by comparison with a standard ammonium sulphate solution similarly Nesslerized.

The last and latest of the indirect methods for the determination of ammonia is the so-called permutit method suggested by Folin and Bell.(1) The reagent which is sold under the trade name of "Permutit" is a complex insoluble sodium salt containing easily replaceable sodium. The chemical basis of the method is the absorption of ammonia by this reagent, its subsequent liberation by alkali and determination by Nesslerization. The accuracy of the method is dependent entirely upon the quality of the reagent, the amount liberated depending upon this variable factor.

1. Folin and Bell. Jr. Biol. Chem. (1917)

oxalate are added, and a strong air current is passed through the mixture until the ammonia has all been removed. The ammonia is collected in ammonia-free water and acid, is neutralized, and the ammonia determined by comparison with a standard ammonia sulphate solution similarly neutralized.

The last and latest of the indirect methods for

the determination of ammonia is the so-called permittivity method suggested by Folin and Bell. (1) The reagent which is sold under the trade name of "Permunit" is a complex insoluble sodium salt containing easily replaceable sodium. The chemical basis of the method is the absorption of ammonia by this reagent, its subsequent liberation by alkali and determination by neutralization. The accuracy of the method is dependent entirely upon the quality of the reagent, the amount liberated depending upon this variable factor.

CARBON DIOXIDE DETERMINATION.

The determination of carbon dioxide as a constituent of the respiratory gas may be accomplished by numerous and diverse procedures. These may be grouped under three heads:

First, the analysis of the gases thru one or another gas analytical procedure.

second, the fixation of the carbon dioxide by some chemical agent and the determination of the material thus formed either by gravimetric, volumetric, colorimetric or thermometric operations,

and third, the optical determination which is dependent upon interference phenomena. Further, each of the above

CARBON DIOXIDE DETERMINATION.

methods proposed, offer a maximum of accuracy with a minimum of simplicity and a minimum of time consumption, while others show the exact opposite of the first.

While our purpose in this work is to secure a method which will offer a sufficient degree of accuracy with a minimum of time consumption, complexity of apparatus and equipment, still in the preliminary work the accurate method must be adopted in order to create a standard of comparison for the less accurate and simpler methods. The following classification will give some idea of the methods available for carbon dioxide determination.

CARROLL DIXON'S DETERMINATION

CARBON DIOXIDE DETERMINATION.

The determination of carbon dioxide as a constituent of the respiratory gases may be accomplished by numerous and diverse procedures. These may be grouped under three heads:

first, the analysis of the gases thro one or another gas analytical procedure;

second, the fixation of the carbon dioxide by some chemical agent and the determination of the material thus formed either by gravimetric, volumetric, colorimetric or thermometric operations,

and third, the optical determination which is dependent upon interference phenomena. Further, each of the first two general types of analyses differ in that certain methods proposed, offer a maximum of accuracy with a minimum of simplicity and a maximum of time consumption, while others show the exact opposite of the first.

While our purpose in this work is to secure a method which will offer a sufficient degree of accuracy with a minimum of time consumption, complexity of apparatus and equipment, still in the preliminary work one accurate method must be adopted in order to create a standard of comparison for the less accurate and simpler methods. The following classification will give some idea of the methods available for carbon dioxide determination.

1. Harriett's determination.

2. Thermometric.

1. Wilcox thermometer.

2. Monroe thermometer.

CARBON DIOXIDE DETERMINATION.

The determination of carbon dioxide as a constituent of the respiratory gases may be accomplished by numerous and diverse procedures. These may be grouped under three heads:

First, the analysis of the gases thro one or another gas analytical procedure;

second, the fixation of the carbon dioxide by some chemical agent and the determination of the material thus formed either by gravimetric, volumetric, colorimetric or thermometric operations,

and third, the optical determination which is dependent upon interference phenomena. Further, each of the first two general types of analyses differ in that certain methods proposed, offer a maximum of accuracy with a minimum of simplicity and a maximum of time consumption, while others show the exact opposite of the first.

While our purpose in this work is to secure a method which will offer a sufficient degree of accuracy with a minimum of time consumption, complexity of apparatus and equipment, still in the preliminary work one accurate method must be adopted in order to create a standard of comparison for the less accurate and simpler methods.

The following classification will give some idea of the methods available for carbon dioxide determination.

I. Determination by Gas Analysis.

Winkler Gas Burette

A. Measurement and Absorption in same burette.

1. Hempel's modified Winkler Burette.
2. Honigman Gas Burette.
3. Bunte Gas Burette.
4. Fridericia collection, measuring and absorption tube.

B. Measurement and Absorption in separate apparatus.

1. Hempel gas apparatus.
2. Elliot gas apparatus.
3. Orsat gas apparatus.
4. Bunsen gas apparatus.
5. Doyere gas apparatus.
6. Herbert and Heim gas apparatus.

C. Measurement and Absorption separate with the addition of means for compensating changes in temperature and barometric pressure.

1. Pettersson gas apparatus.
2. Hempel gas apparatus.
3. Pettersson and Palmqvist gas apparatus.
4. Haldane gas apparatus.
5. Henderson gas apparatus.

D. Change of density with changing chemical composition.

1. Nernst Gas Balance.

II. Determination after the Fixation of the Carbon Dioxide.

A. Gravimetric.

1. Haldane Carbon dioxide absorber.
2. Benedict " " "

B. Volumetric.

1. Saussure and Pettenkofer titration.
2. Hesse titration.
3. Henderson titration.

C. Colorimetric.

1. Marriott determination.

D. Thermometric.

1. Wilcox thermometer.
2. Munroe thermometer.

I. Determination by Gas Analysis.

Winkler Gas Burette

A. Measurement and Absorption in same burette.

1. Hempel's modified Winkler burette.
2. Hönigsmann Gas Burette.
3. Bunte Gas Burette.
4. Fiske's collection, measuring and absorption tube.

B. Measurement and Absorption in separate apparatus.

1. Hempel gas apparatus.
2. Elliot gas apparatus.
3. Orsat gas apparatus.
4. Smith gas apparatus.
5. Coyne gas apparatus.
6. Herbert and Reid gas apparatus.

C. Measurement and Absorption separate with the addition of means for compensating changes in temperature and barometric pressure.

1. Pettersson gas apparatus.
2. Hempel gas apparatus.
3. Pettersson and Palmqvist gas apparatus.
4. Hallane gas apparatus.
5. Henderson gas apparatus.

D. Change of density with changing chemical composition.

1. Marsh Gas Balance.

II. Determination after the fixation of the Carbon Dioxide.

A. Gravimetric.

1. Hallane Carbon Dioxide absorber.
2. Benedict "

B. Volumetric.

1. Sauerbre and Bettendorfer titration.
2. Masse titration.
3. Henderson titration.

C. Colorimetric.

1. Harriott determination.

D. Thermometric.

1. Wilcox thermometer.
2. Hunsche thermometer.

III. Determination with interferometer.

The various types of methods will be considered in order of classification with an estimation of their applicability to the determination of alveolar carbon dioxide, and of their availability for use in the present problem. The methods for the collection of the alveolar air, which is of course pre-requisite to all procedures for carbon dioxide determination involved in this study, are an important phase of the subject and will be subsequently considered. For the time being we will consider that the gas is available for the determination, in one manner or another, of its carbon dioxide content.

Measurement of the amount of the gas is the first operation in the use of a gas analytical method. Such measurement is best accomplished by the determination of the volume of the gas under known conditions of temperature, of pressure and of tension of liquids that are present. The earliest and simplest apparatus for gas measurement is the Winkler Gas Burette (1), commonly called the Hempel Gas Burette. As a matter of fact, however, Hempel's original burette is a modification of the Winkler. This Winkler Burette consists of two tubes A and B which are set in iron feet. Inside the feet the tubes are bent at right angles and conically drawn out, the ends projecting from the feet. The two tubes are connected by rubber tubing slipped over these projections. The measuring tube A ends at the top in a capillary tube over which a short piece of rubber tubing closed with a pinchcock

is slipped. A contains between the base of the capillary and a graduation a little above the iron foot one hundred cubic centimeters and is graduated to fifths of a cubic centimeter. B is used as a leveling tube and as a means of introducing absorbing solutions into the measuring burette. Manipulation of this first apparatus must have been something of an ordeal, and would require a high degree of technique to secure results even approximating those obtained by present day methods.

The procedure was as follows:

Fill the tubes A and B with water, taking care to drive all air out of the connecting rubber tube by raising or lowering the tubes. Then, join the burette to the vessel containing the gas by means of a glass or rubber tube filled with water. To fill the burette with the gas to be examined, close the connecting tube between A and B and pour the water out of B. Place the leveling tube on the floor and open the pinchcock at the top of A. The water will flow into the leveling-tube and the gas will be drawn into the burette. When A is filled with gas close the pinchcock, disconnect from the gas-holder and after the liquid has run down from the walls of the burette, take up the tubes by the iron feet and by raising or lowering bring the water in the tubes to the same level. The gas is now under atmospheric pressure, and its volume is read off. To bring as much as possible of the absorbent into the burette, lower the leveling-tube until the expanded gas begins

is slipped. A container between the base of the capillary and a graduation a little above the iron feet are marked with a centimeter and is graduated to fifths of a centimeter. It is used as a leveling tube and as a means of introducing absorbing solutions into the measuring burette. Manipulation of this first apparatus must have been something of an ordeal, and would require a high degree of technique to secure results even approximating those obtained by present day methods.

The procedure was as follows:

Fill the tubes A and B with water, taking care to drive all air out of the connecting rubber tube by raising or lowering the tubes. Then, join the burette to the vessel containing the gas by means of a glass or rubber tube filled with water. To fill the burette with the gas to be examined, close the connecting tube between A and B and pour the water out of B. Place the leveling tube on the floor and open the pinchcock at the top of A. The water will flow into the leveling-tube and the gas will be drawn into the burette. When A is filled with gas close the pinchcock, disconnect from the gas-holder and after the liquid has run down from the walls of the burette, take up the tubes by the iron feet and by raising or lowering bring the water in the tubes to the same level. The gas is now under atmospheric pressure, and its volume is read off. To bring as much as possible of the absorbent into the burette, lower the leveling-tube until the expanded gas begins

to enter the rubber tube, close the connecting tube and pour the greater part of the water out of B. The absorbing liquid is now poured into B and the tube raised as far as the rubber tube permits; in this way a considerable amount of the reagent, diluted with the water in the rubber tube is brought into the burette. Closing the tube at the base of A, bring the gas into close contact with the absorbent by vigorously shaking the burette. When no further decrease of the volume of the enclosed gas takes place, the reading is made as before. The difference in volume gives the amount of the absorbed gas.

Because of its solubility in water, the initial volume of a gas containing carbon dioxide cannot be made over water. Hempel therefore modified the original Winkler Burette. He added a three way cock at the bottom of the measuring burette. The burette is filled by drawing the gas thro the tube by means of suction applied at the lower cock and then closing both ends. Atmospheric pressure is obtained by opening the upper end momentarily. The absorbing liquids are introduced by means of a funnel connected with the longitudinal opening of the three way cock, by a piece of rubber tubing. Except for this the manipulation is the same as is described for the simple Winkler Burette.

This method of Winkler and its modification by Hempel are the basis not only of gas analysis involving separate absorption pipettes, as the various methods of Hempel, Orsat and their followers, but also of the Honigman, Bunte and Fridericia methods in which, as in the Winkler, the absorption takes place within the measuring

burette. These latter methods we will consider first.

Honigmann (1) did away with the leveling tube entirely, using for his determination simply a one hundred cubic centimeter burette closed at the top with a glass stopcock, the lower end being drawn out to smaller diameter to permit of a piece of rubber tubing being easily slipped over it. The absorbing liquid is placed in a very tall glass cylinder. The gas to be analyzed is passed through the burette until all the air previously contained has been displaced. The stopcock is then closed and the rubber tube is immersed in a solution of potassium hydroxide. The burette is then lowered into the solution until the surrounding liquid stands at the zero point, the lowest graduation upon the tube, the stopcock is opened and the liquid allowed to rise inside the burette to the same mark. The tube now contains one hundred cubic centimeters of the gas at atmospheric pressure. The absorption of the gas is effected by inverting the burette and allowing the alkali to flow along the walls of the burette. The end of the rubber tube must of course remain below the surface of the liquid in the cylinder. After absorption the burette is lowered in the solution until the liquid surfaces within and without stand at the same height. The percentage of carbon dioxide present is read directly. The Honigmann burette is suited only to a rapid approximate determination of carbon dioxide in gas mixtures which contain fairly high percentages of that constituent.

burette. These latter methods we will consider first.
 Hönigsmann (1) did away with the leveling tube
 entirely, using for his determination simply a one hundred
 cubic centimeter burette closed at the top with a glass
 stopcock, the lower end being drawn out to smaller diam-
 eter to permit of a piece of rubber tubing being easily
 slipped over it. The absorbing liquid is placed in a very
 tall glass cylinder. The gas to be analyzed is passed into
 the burette until all the air previously contained has been
 displaced. The stopcock is then closed and the rubber tube
 is immersed in a solution of potassium hydroxide. The bu-
 rette is then lowered into the solution until the surround-
 ing liquid stands at the zero point, the lowest graduation
 upon the tube, the stopcock is opened and the liquid allowed
 to rise inside the burette to the same mark. The tube now
 contains one hundred cubic centimeters of the gas at atmos-
 pheric pressure. The absorption of the gas is effected by
 inverting the burette and allowing the alkali to flow along
 the walls of the burette. The end of the rubber tube must
 of course remain below the surface of the liquid in the
 cylinder. After absorption the burette is lowered in the
 solution until the liquid surfaces within and without stand
 at the same height. The percentage of carbon dioxide present
 is read directly. The Hönigsmann burette is suited only to
 a rapid approximate determination of carbon dioxide in gas
 mixtures which contain fairly high percentages of that con-
 stituent.

The Bunte (1) apparatus consists of a burette, a suction bottle and a beaker of alkali. The burette is bulbed to avoid extreme length and is provided at the top with a funnel and a three-way cock and at the bottom with an ordinary cock. The gas is collected over water as in the original Winkler, the suction bottle taking the place of the leveling tube. To introduce the reagent the lower tip is immersed in the alkali and the lower cock opened. After the absorption of the carbon dioxide the reagent is washed from the burette by water introduced thro the funnel, the tip of the burette being all the time under water. Water then rises in the burette, filling the vacuum created by the removal of the carbon dioxide and the reading is made. The difference between the first and second readings represents the number of cubic centimeters of carbon dioxide present. Readings are made with the funnel filled to the same level and the upper cock open; the pressure being thus atmospheric pressure plus the pressure of the height of the column of water in the funnel. Determinations made with this apparatus are of necessity inaccurate for the gas is brought into contact with large volumes of water which are unsaturated with the gas mixture and will, therefore, absorb some of the gas of the sample. Reagent is also wasted because of its dilution with wash water. The use of the burette is like the Honigmann; limited to the approximate determination of carbon dioxide.

The Bunte (I) apparatus consists of a burette, a suction bottle and a beaker of alkali. The burette is dipped to avoid extreme length and is provided at the top with a funnel and a three-way cock and at the bottom with an ordinary cock. The gas is collected over water as in the original Winkler, the suction bottle taking the place of the leveling tube. To introduce the reagent the lower tip is immersed in the alkali and the lower cock opened. After the absorption of the carbon dioxide the reagent is washed from the burette by water introduced into the funnel, the tip of the burette being all the time under water. Water then rises in the burette, filling the vacuum created by the removal of the carbon dioxide and the reading is made. The difference between the first and second readings represents the number of cubic centimeters of carbon dioxide present. Readings are made with the funnel filled to the same level and the upper cock open; the pressure being thus atmospheric pressure plus the pressure of the height of the column of water in the funnel. Determinations made with this apparatus are of necessity inaccurate for the gas is brought into contact with large volumes of water which are unsaturated with the gas mixture and will, therefore, absorb some of the gas of the sample. Reagent is also wasted because of its dilution with wash water. The use of the burette is like the Hönigsmann; limited to the approximate determination of carbon dioxide.

The most recent development of the Winkler Gas Burette along this line where the absorption and measurement are made in one tube, is the Fridericia (1) tube. This Fridericia tube and the method of its use are particularly interesting to us in this study, for it was devised by Fridericia and has been recommended by many others as a suitable method for the clinical determination of alveolar carbon dioxide. The apparatus necessary for the determination consists of the Fridericia absorbing pipette, in which the air is collected and analyzed, a cylinder of water at room temperature in which the tube may be immersed, and a five percent solution of caustic soda. The absorbing pipette is essentially a U-shaped tube with stopcocks so arranged, that it may be opened from end to end, allowing the air to be blown through, or closed so that one chamber of exactly one hundred cubic centimeters capacity may be closed off from the rest of the tube. The lower section of this chamber is graduated in tenths of one percent of the total volume of the chamber.

The method of procedure is as follows:
At the end of a normal inspiration the subject is asked to blow through the open tube, emptying the lungs as far as possible. The upper end of the measured portion of the tube is closed and the air is cooled to room temperature by immersion in the tank, contraction within the measuring space being compensated by air from the open arm, the assumption being made that its composition is similar to that within the measuring space. How far diffusion vitiates this assumption has not been determined. Alkali is introduced into the

The most recent development of the Winkler Gas
method along this line is the absorption and measure-
ment made in the tube, in the Winkler (1) tube.
This Winkler tube and the method of its use are partic-
ularly interesting to us in this study, for it was devised
by Winkler and has been recommended by many others as
a reliable method for the chemical determination of dissolved
carbon dioxide. The apparatus necessary for the determination
consists of the Winkler absorbing pipette, in which the
air is collected and analyzed, a cylinder of water of room
temperature in which the tube may be immersed, and a five
percent solution of caustic soda. The absorbing pipette is
essentially a U-shaped tube with reservoirs at each end,
that it may be opened from end to end, allowing the air to
be drawn through, or closed so that one chamber of exactly
one hundred cubic centimeters capacity may be closed off
from the rest of the tube. The lower section of this cham-
ber is graduated in terms of one percent of the total volume
of the chamber.

The method of procedure is as follows:

At the end of a normal inspiration the subject is asked to
breathe through the open tube, emptying the lungs as far as
possible. The upper end of the measured portion of the tube
is closed and the air is cooled to room temperature by im-
mersion in the tank, connection within the measuring space
being compensated by air from the open end. The assumption
being made that the composition is similar to that within
the measuring space. Now the diffusion within this space
has not been determined. Aikoff is introduced into the

air chamber and the carbon dioxide is absorbed by it. The tube is returned to the tank, the lower cock is opened, and the carbon dioxide containing alkali is allowed to drain out; the water then rises in the tube, filling the vacuum created by the removal of the carbon dioxide. The percentage of carbon dioxide is read directly. This percentage as any of the percentages obtained by a gas analysis procedure may be changed to millimeters of mercury pressure in dry air by multiplying the difference between barometric pressure at the time of the test and the tension of aqueous vapor at 37.5°C which is 48 mm. mercury. The advantages of this method include its simplicity of design and of technique required for its use combined with the possibility of easy transportation of the apparatus to the bedside.

Because of the difficulties of manipulation and the need of cleaning the apparatus between each determination where the measurement and absorption is carried out in the same tube, the original gas burette was further modified by the addition of separate absorption apparatus. These were connected with the measuring burette by capillary tubes of as short length as possible, the air being forced over into the absorbers by the raising and lowering of the leveling tube. The error caused by measurement over water, the carbon dioxide being very soluble in water, was obviated by the use of mercury in the leveling tube. A drop of water placed in the measuring burette insures saturation of the gas. Further, the measuring burette was immersed in a water jacket thereby preventing changes in the gas volume due to

air chamber and the carbon dioxide is absorbed by it. The tube is returned to the tank, the lower cock is opened, and the carbon dioxide containing liquid is allowed to drain out; the water then rises in the tube, filling the vacuum created by the removal of the carbon dioxide. The percentage of carbon dioxide is read directly. This percentage as any of the percentages obtained by a gas analysis procedure may be changed to millimeters of mercury pressure in dry air by multiplying the difference between barometric pressure at the time of the test and the tension of aqueous vapor at 37.5°C which is 48 mm. mercury. The advantages of this method include the simplicity of design and of technique required for its use combined with the possibility of easy transportation of the apparatus to the bedside.

Because of the difficulties of manipulation and the need of cleaning the apparatus between each determination where the measurement and absorption is carried out in the same tube, the original gas burette was further modified by the addition of separate absorption apparatus. These were connected with the measuring burette by capillary tubes of as short length as possible, the air being forced over into the absorbers by the raising and lowering of the leveling tube. The error caused by measurement over water, the carbon dioxide being very soluble in water, was avoided by the use of mercury in the leveling tube. A drop of water placed in the measuring burette insures saturation of the gas. Further, the measuring burette was immersed in a water jacket thereby preventing changes in the gas volume due to

variations in temperature. The types of apparatus designed by Hempel (1), Elliott (2), Orsat (3) and many others show one or all of these modifications.

The third division in the classification of gas analytical methods, contains the latest descendants of the original Winkler burette. Their most important common feature is the use of a tube so connected with the measuring burette as to compensate for changes in atmospheric pressure. The possibility of such compensation was first suggested by Pettersson. The later apparatus of Hempel (4), the Pettersson-Palmqvist (5) apparatus and types designed by Haldane (6) in England and Henderson (7) and his co-workers in this country, all make use of a compensation tube. The Haldane portable apparatus offers a means of exact gas analysis, as is amply attested by the work of the many investigators making use of it. It has been adopted as the standard of comparison in this work, and description of it and the technique of its manipulation will serve to illustrate the methods employed in modern gas analysis.

The apparatus and method of use are fully described by Haldane in his brief treatise on "Methods of Air Analysis". In brief the procedure is as follows: The gas to be analyzed is drawn into a measuring burette

- | | | | |
|----------|----------------|----------------------|-----------|
| 1 and 4. | Denis | Gas Analysis. | (1913) |
| 2. | Elliott Scott, | Standard Methods of | |
| | | Chemical Analysis. | |
| 3. | Orsat. | Chemical News. | (1874) |
| 5. | Pettersson. | Zeit. f. Anal. Chem. | (1886) |
| 6. | Haldane. | Methods of Gas Anal. | (1912) |
| 7. | Henderson. | Jr. Biol. Chem. | 38 (1918) |

variations in temperature. The types of apparatus designed by Hempel (1), Elliott (2), Orsat (3) and many others show one or all of these modifications.

The third division in the classification of gas analytical methods, contains the latest developments of the original Winkler burette. Their most important common feature is the use of a tube so connected with the measuring burette as to compensate for changes in atmospheric pressure. The possibility of such compensation was first suggested by Patterson. The later apparatus of Hempel (4), the Patterson-Balsqvist (5) apparatus and types designed by Haldane (6) in England and Henderson (7) and his co-workers in this country, all make use of a compensation tube. The Haldane portable apparatus offers a means of exact gas analysis, as is amply attested by the work of the many investigators making use of it. It has been adopted as the standard of comparison in this work, and description of it and the technique of its manipulation will serve to illustrate the methods employed in modern gas analysis.

The apparatus and method of use are fully described by Haldane in his brief treatise on "Methods of Air Analysis". In brief the procedure is as follows:

The gas to be analyzed is drawn into a measuring burette

1 and 4.	Copie	Gas Analysis.	(1913)
2.	Elliott Scott.	Standard Methods of Chemical Analysis.	(1913)
3.	Orsat.	Chemical News.	(1874)
5.	Patterson.	Zeit. f. Anal. Chem.	(1888)
6.	Haldane.	Methods of Gas Anal.	(1913)
7.	Henderson.	Pr. Biol. Chem.	(1913)

surrounded by a water jacket and is there saturated with water over mercury, cooled to room temperature and measured. In the same water jacket is a control tube of about the same volume as the burette. The measuring tube and the control tube are connected through the potash absorption pipette which serves as a pressure guage. Before every reading of the burette the potash is brought to the same level. The air in the burette is freed from carbon dioxide by forcing it back and forth into the absorption chamber containing potassium hydroxide. Measurement of the air is made before and after the operation and the percentage of carbon dioxide calculated. The method is one which gives a very high degree of accuracy with a minimum of time consumption and requires only a fair degree of technical skill.

The fourth group of methods, namely that depending upon varying densities of gas mixtures (1) with changing composition such as the gas balance, are all continuous procedures and all wholly unsuited for intermittent analysis. As but one phase of the respiratory cycle is subjected to analysis, only static methods are applicable.

Gas analysis is not, however, the only means available for the determination of this gas. Gravimetric methods are accurate and valuable where the determination of large quantities of the gas is desired. It is the standard method for use with respiration calorimeters (2). The carbon dioxide is absorbed by granulated soda lime and the carbon dioxide determined by the difference in the

1. Nernst and Hunter. Zeit. für Physikal. Chem. (1905)
2. Carpenter. Carnegie Instit. Public 216 (1915)

surrounded by a water jacket and is there saturated with water over mercury, cooled to room temperature and massed. In the same water jacket is a control tube of about the same volume as the burette. The measuring tube and the control tube are connected through the potash absorption pipette which serves as a pressure gauge. Before every reading of the burette the potash is brought to the same level. The air in the burette is freed from carbon dioxide by forcing it back and forth into the absorption chamber containing potassium hydroxide. Measurement of the air is made before and after the operation and the percentage of carbon dioxide calculated. The method is one which gives a very high degree of accuracy with a minimum of time consumption and requires only a fair degree of technical skill.

The fourth group of methods, namely that depending upon varying densities of gas mixtures (1) with changing composition such as gas balance, are all continuous procedures and are wholly unsuited for intermittent analysis. As but one phase of the respiratory cycle is subjected to analysis, only static methods are applicable. Gas analysis is not, however, the only means available for the determination of this gas. Gravimetric methods are accurate and valuable where the determination of large quantities of the gas is desired. It is the standard method for use with respiration calorimeters (2). The carbon dioxide is absorbed by granulated soda lime and the carbon dioxide determined by the difference in the

weight of the soda lime before and after absorption.

Such a method requires of course the removal of all water from the air before it enters the carbon dioxide absorber. The absorber consists generally of either a metal cylinder or a glass container filled with the soda lime and provided with air tight connections permitting of the introduction and withdrawal of the air. Such a method might conceivably be used for this work. The subject could breathe for a period of ten minutes or so through a system containing a small meter for measuring the air, a sulphuric acid container for removing the water and a carbon dioxide absorber. The carbon dioxide tension could then be determined by weighing the absorber, the amount being divided by the volume of expired air. The oxygen level is maintained by use of a tank of pure gas in controlled connection. Such apparatus as the Benedict portable respiration unit,⁽¹⁾ is designed for this and kindred purposes. The relatively high cost of an installation, together with the somewhat complicated manipulation precludes the general use of this apparatus and limits it to institutions where these conditions can be met.

Another possibility is the fixation of carbon dioxide by some liquid reagent and its titrimetric determination. The procedures available are those of Pettenkofer⁽²⁾ Henderson⁽³⁾ and Marriott⁽⁴⁾. The first two are essentially the same. A measured volume of air is in each case treated

- | | | | |
|-----------------|--------------------------|-----|--------|
| 1. Carpenter. | Carnegie Instit. Public. | 216 | (1915) |
| 2. Pettenkofer. | Am. Jr. Phys. | | (1912) |
| 3. Henderson. | " " " | | (1912) |
| 4. Marriott. | Jr. Am. Med. Association | | (1916) |

weight of the soda lime before and after absorption. Such a method requires of course the removal of all water from the air before it enters the carbon dioxide absorber. The absorber consists generally of either a metal cylinder or a glass container filled with the soda lime and provided with air tight connections permitting of the introduction and withdrawal of the air. Such a method might conceivably be used for this work. The subject could breathe for a period of ten minutes or so through a system containing a small meter for measuring the air, a sulphuric acid container for removing the water and a carbon dioxide absorber. The carbon dioxide tension could then be determined by weighing the absorber, the amount being divided by the volume of expired air. The oxygen level is maintained by use of a tank of pure gas in controlled connection. Such apparatus as the Benedict portable respiration unit, is designed for this and related purposes. The relatively high cost of an installation, together with the somewhat complicated manipulation precludes the general use of this apparatus and limits it to installations where these conditions can be met. Another possibility is the fixation of carbon dioxide by some liquid reagent and the titrimetric determination. The procedures available are those of Pettenkofer (2) Henderson (3) and Harriott (4). The first two are essentially the same. A measured volume of air is in each case treated

1. Carpenter.	215	(1915)
2. Pettenkofer.	Am. J. Hyg.	(1915)
3. Henderson.	" "	(1915)
4. Harriott.	Br. Med. Association	(1915)

with an excess of barium hydroxide solution which quantitatively absorbs carbon dioxide forming insoluble barium carbonate. Phenolphthalein is added to the solution and the excess of barium hydroxide is titrated with hydrochloric acid. From the amount of alkali used to absorb the carbon dioxide the amount of the latter is calculated. Pettenkofer uses a flask of known capacity and fills it with the air to be analyzed by means of bellows, notes the pressure and temperature, adds by means of a pipette one hundred cubic centimeters of barium hydroxide solution, shakes vigorously, pours into a dry flask, and pipettes out twenty-five cubic centimeters which he titrates with the hydrochloric acid. The strength of the barium hydroxide in terms of acid being known, the amount and percent of carbon dioxide can be calculated at 0°centigrade and 760 millimeters of pressure. The method of Henderson is much simpler and unquestionably more exact. His apparatus permits of the air being sucked directly into a measured pipette from the Haldane gas collection tube, where it is treated with the barium hydroxide. The alkali is then withdrawn into a flask, the precipitate allowed to settle and the titration carried out with hydrochloric acid. The value from the clinical standpoint of both of these methods is lessened by the fact that standardized alkali and acid are necessary for the determination.

The determination by the method described by Marriott while likewise dependent upon the hydrogen ion concentration varies widely from the two just described. The apparatus consists of eight test tubes, containing standard

with an excess of barium hydroxide solution which gradually
 tatively absorbs carbon dioxide forming insoluble barium
 carbonate. Phenolphthalein is added to the solution and
 the excess of barium hydroxide is titrated with hydrochloric
 acid. From the amount of alkali used to absorb the carbon
 dioxide the amount of the latter is calculated. Potentiometer
 uses a flask of known capacity and fills it with the air to
 be analysed by means of bellows, notes the pressure and tem-
 perature, adds by means of a pipette one hundred cubic cen-
 timeters of barium hydroxide solution, shakes vigorously,
 pours into a dry flask, and pipettes out twenty-five cubic
 centimeters which is titrated with the hydrochloric acid.
 The strength of the barium hydroxide in terms of acid being
 known, the amount and percent of carbon dioxide can be cal-
 culated at 0° centigrade and 760 millimeters of pressure.
 The method of Henderson is much simpler and unexceptionally
 more exact. His apparatus permits of the air being sucked
 directly into a measured pipette from the Euland gas col-
 lection tube, where it is treated with the barium hydroxide.
 The alkali is then withdrawn into a flask, the precipitate
 allowed to settle and the titration carried out with hydro-
 chloric acid. The value from the clinical standpoint of
 both of these methods is lessened by the fact that standard-
 ized alkali and acid are necessary for the determination.
 The determination by the method described by
 Marshall while likewise dependent upon the hydrogen ion con-
 centration varies widely from the two just described. The
 apparatus consists of eight test tubes, containing standard

phosphate solutions, a standard bicarbonate solution, a small test tube, a glass tube drawn out to a capillary point and a box for color comparison. The method depends upon the fact that if a current of air containing carbon dioxide is passed through a solution of sodium carbonate or bicarbonate until the solution is saturated, the final solution will contain sodium bicarbonate and dissolved carbon dioxide. The reaction of such a solution will depend upon the relative amounts of the alkaline bicarbonate and the acid carbon dioxide present. This, in turn, will depend on the tension of carbon dioxide in the air with which the mixture was saturated, and will be independent of the volume of air blown through it, provided saturation has been obtained. High tensions of carbon dioxide change the reaction of the solution toward the acid side. Low tensions have the reverse effect; hence, the reaction of such a solution is a measure of the tension of carbon dioxide in the air with which it has been saturated. The reaction of such a solution may be determined by adding to it an indicator such as phenolsulphone-phthalein which shows over a considerable range of reaction definite color changes. A certain color indicates a certain reaction. Solutions of a given reaction may be prepared by mixing acid and alkaline phosphates in definite proportions. Such solutions according to Marriott may be kept unaltered for long periods of time and can be used as standards for comparison. The eight standards are so prepared as to show a hydrogen ion concentration equivalent to that shown by carbon dioxide tensions of 10,15,20,25,30,35,40 and 45 mili-

phosphate solutions, a standard bicarbonate solution, a small test tube, a glass tube drawn out to a capillary point and a box for color comparison. The method depends upon the fact that if a current of air containing carbon dioxide is passed through a solution of sodium carbonate or bicarbonate until the solution is saturated, the final solution will contain sodium bicarbonate and dissolved carbon dioxide. The reaction of such a solution will depend upon the relative amounts of the alkaline bicarbonates and the acid carbon dioxide present. This, in turn, will depend on the tension of carbon dioxide in the air with which the mixture was saturated, and will be independent of the volume of air blown through it, provided saturation has been obtained. High tensions of carbon dioxide change the reaction of the solution toward the acid side. Low tensions have the reverse effect; hence, the reaction of such a solution is a measure of the tension of carbon dioxide in the air with which it has been saturated. The reaction of such a solution may be determined by adding to it an indicator such as phenolphthalein which shows over a considerable range of reaction definite color changes. A certain color indicates a certain reaction. Solutions of a given reaction may be prepared by mixing acid and alkaline phosphates in definite proportions. Such solutions according to Harriott may be kept unaltered for long periods of time and can be used as standards for comparison. The eight standards are so prepared as to show a hydrogen ion concentration equivalent to that shown by carbon dioxide tensions of 10, 15, 20, 25, 30, 35, 40 and 45 mm.

meters of mercury in saturated bicarbonate solution. Comparisons are made with a flat black metal box, backed with opal glass and containing holes of the standard and the unknown. For the analysis of a sample of air about two or three cubic centimeters of the standard bicarbonate solution are poured into a clean test tube of the same diameter as the standard tubes. Air collected in a rubber bag is then blown through the solution, by means of a glass tube drawn out to a capillary point, until the solution is saturated as shown by the fact that no further color change occurs. The tube is stoppered and the color immediately compared with that in the standard tubes. The method is a very rapid one, its value is dependent upon the ease or difficulty with which correct standards may be prepared, the length of time that they will maintain their integrity, and the ability of the operator to apprehend fine color distinctions.

While according to Marriott the standards for comparison are easily prepared and will retain their integrity for a long time, work which has already been done in connection with this research would indicate that such an assertion betrays an unwarranted degree of optimism on the part of its author. Marriott's directions for the preparation of the standards are as follows:

"Fifteenth molecular acid potassium phosphate:

Of the pure recrystallized salt (KH_2PO_4), 9.078 gms. are dissolved in distilled water, 200cc of 0.01 percent phenol-sulphonaphthalein solution are added, and the whole is made

meters of mercury in saturated bicarbonate solution. Com-
parisons are made with a flat black metal box, backed with
opal glass and containing pieces of the standard and the
unknown. For the analysis of a sample of air about two
or three cubic centimeters of the standard bicarbonate
solution are poured into a clean test tube of the same li-
quor as the standard tubes. Air collected in a rubber
bag is then blown through the solution, by means of a
glass tube drawn out to a capillary point, until the so-
lution is saturated as shown by the fact that no further
color change occurs. The tube is stoppered and the color
immediately compared with that in the standard tubes. The
method is a very rapid one, its value is dependent upon
the ease or difficulty with which correct standards may be
prepared, the length of time that they will maintain their
integrity, and the ability of the operator to apprehend fine
color distinctions.

While according to Merritt the standards for com-
parison are easily prepared and will retain their integrity
for a long time, work which has already been done in con-
nection with this research would indicate that such an asser-
tion betrays an unwarranted degree of optimism on the part
of its author. Merritt's directions for the preparation
of the standards are as follows:

"Fifty cubic centimeters of a solution of potassium phosphate:
Of the pure reagent grade K_2HPO_4 , 5.075 gms. are
dissolved in distilled water, 300cc of 0.01 percent phenol-
anthracene solution are added, and the whole is made

up to one liter.

Fifteenth molecular alkaline sodium phosphate:

The pure recrystallized salt ($\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$) is exposed to the air for from ten days to two weeks, protected from the dust. Ten molecules of water of crystallization are given off and a salt of the formula $\text{Na}_2\text{HPO}_4 \cdot 2\text{H}_2\text{O}$ is obtained. Of this salt 11.876 gms. are dissolved in water, 200cc of 0.01 percent. phenolsulphonephthalein solution are added, and the whole is made up to one liter.

Proportions in which solutions are mixed:

mm.	10	15	20	25	30	35	40	45
Potassium phosphate	17.8	25.2	31.0	35.7	40.5	45.0	47.0	50.2
Sodium phosphate	82.2	74.8	69.0	64.3	59.5	55.0	53.0	49.8

The standard alkali solution for absorption is prepared either by weighing out 0.530 gms. of dessicated sodium carbonate or by measuring accurately 100cc of tenth normal sodium hydroxide into a one liter volumetric flask. 200cc of 0.01 percent. phenolsulphonephthalein are added, and the whole is made up to the mark with distilled water. Carbon dioxide from a cylinder or from the lungs may be passed through this solution to convert the alkali into bicarbonate or the solution may simply be used as it is, as the alveolar air that will be blown through the solution subsequently will accomplish the same purpose.

Owing to the difficulty of estimating the amount of water of crystallization in the sodium phosphate salt

up to one liter.

Filtered solution alkaline sodium phosphate:

The pure recrystallized salt ($\text{Na}_2\text{HPO}_4 \cdot 12\text{H}_2\text{O}$) is exposed

to the air for from ten days to two weeks, protected

from the heat. Ten volumes of water of crystallization

are given off and a salt of the formula Na_2HPO_4 is

obtained. Of this salt 11.875 gms. are dissolved in

water, 300cc of 0.01 percent. phenolphthalein so-

lution are added, and the whole is made up to one liter.

Proportions in which solutions are mixed:

mm.	10	15	20	25	30	35	40	45
Potassium phosphate	17.8	25.2	31.9	38.7	40.5	45.0	47.0	50.2
Sodium phosphate	32.2	24.8	22.0	24.3	29.5	28.0	23.0	19.8

The standard alkali solution for absorption is

prepared either by weighing out 0.550 gms. of desiccated

sodium carbonate or by measuring accurately 100cc of tenth

normal sodium hydroxide into a one liter volumetric flask.

300cc of 0.01 percent. phenolphthalein are added,

and the whole is made up to the mark with distilled water.

Carbon dioxide from a cylinder or from the lungs may be

passed through this solution to convert the alkali into

bicarbonate or the solution may simply be used as it is.

As the alveolar air that will be blown through the solution

subsequently will accomplish the same purpose.

Owing to the difficulty of estimating the amount

of water of crystallization in the sodium phosphate salt

the preparation of a fifteenth molecular solution is not the simple matter that it seems. Standards prepared according to Marriott's direction have not given solutions that check with analyses made by the Haldane apparatus nor with each other. For the purposes of this research fifteenth molecular mono- and di-sodium phosphate solutions are prepared from molar solutions of pure phosphoric acid and carbonate free sodium hydroxide. These are then mixed in the proportions indicated above. Since the action of the alkali upon the glass of the tubes even in a short time causes a change in the hydrogen ion concentration of the standards, the solutions thus prepared are placed in paraffin lined delivery bottles from which the standard tubes may be filled as often as is necessary. These tubes are closed with rubber stoppers during the period of use.

Thermoscopic (1) (2) determinations of carbon dioxide are possible but save for the noting of such possibility are of little interest to us here. The method has not been applied to biological determinations. It depends upon the measurement of the rise of temperature of pulverized sodium hydroxide over which the air containing the carbon dioxide is passed. The percentage of carbon dioxide is registered directly on the Thermoscope scale.

Widely at variance with all methods previously described, but one which for institutional work at any rate, might prove of real value, is the determination by the use of the interferometer (3). The principle of the

- | | | |
|---------------------------------------|--------|--------|
| 1. Engineering | 95 787 | (1913) |
| 2. Am. Gas Light Journal | 99 62 | (1913) |
| 3. Hahn and Heim. Berl. Klin. Wochen. | | (1913) |

The preparation of a lithium molecular solution is not the simple matter that it seems. It has been prepared according to Kretschmer's direction have not given solutions that check with analyses made by the Helium apparatus nor with each other. For the purposes of this research lithium molecular and lithium phosphate solutions are separated from other solutions of pure phosphoric acid and carbonate free sodium hydroxide. These are then mixed in the proportions indicated above. Since the action of the alkali upon the glass of the tubes even in a short time causes a change in the hydrogen ion concentration of the standard, the solutions have prepared are placed in paraffin lined delivery bottles from which the standard tubes may be filled as often as is necessary. These tubes are closed with rubber stoppers during the period of use.

Thermographic (1) (2) Determinations of carbon dioxide are possible but have for the making of such possibility are of little interest to us here. The method has not been applied to biological determinations. It depends upon the measurement of the rise of temperature of oxygenated sodium hydroxide over which the air containing the carbon dioxide is passed. The percentage of carbon dioxide is registered directly on the thermographic scale.

Widely as variance with all methods previously described, but one which for institutional work at any rate, might prove of real value, is the determination by the use of the interferometer (3). The principle of the

1. Engineering (1913) 95 757
2. Am. Gas Light Journal (1913) 93 62
3. Ehm and Helm. Berl. Klin. Wochens. (1913)

interferometer in its application to the composition of gases may be described briefly as follows: First, a simple optical device permits the passage of selected beams of light to a screen and the interference phenomena rendered visible. Where crest meets crest in successive superimposed waves of light, there is an intensification of the illumination; where crest meets trough there is complete interference and consequent extinction. Naturally, all gradations between these limits are possible. In the interferometer in question, the apparatus is constructed in duplicate so that two interference pictures are thrown on the screen, one directly above the other. The light passes through two small chambers, one associated with each spectrum. So long as the index of refraction of the medium filling these two chambers is identical, the pictures of the upper and lower are directly comparable. If, however, the index of refraction of one medium is altered, there is a shifting of the lines to the right or left, as the case may be. By a simple screw device carrying a scale and vernier, this shift can be measured and thus differences in refraction index ascertained. To apply this to carbon dioxide determinations in alveolar air, it is only necessary to calibrate the instrument, using pure carbon dioxide-free air as the standard, and then adding to it successive portions of carbon dioxide, placing this second gas mixture in the lower comparison chamber, the pure air filling the upper, and then measur-

interferometer in its application to the composition

of gases may be described briefly as follows:

First, a simple optical device permits the passage of selected beams of light to a screen and the interference phenomena rendered visible. Where great masses of successive superimposed waves of light, there is an intensification of the illumination; where great masses travel there is complete interference and consequent extinction. Naturally, all variations between these limits are possible. In the interferometer in question, the apparatus is constructed in duplicate so that two interference pictures are thrown on the screen, one directly above the other. The light passes through two small chambers, one associated with each spectrum. So long as the index of refraction of the medium filling these two chambers is identical, the pictures of the upper and lower are directly comparable. If, however, the index of refraction of one medium is altered, there is a shifting of the lines to the right or left, as the case may be. By a simple screw device carrying a scale and vernier, this shift can be measured and thus differences in refraction index ascertained. To apply this to carbon dioxide determinations in a vessel of air, it is only necessary to calibrate the instrument, using pure carbon dioxide-free air as the standard, and then adding to it successive portions of carbon dioxide, placing this second gas mixture in the lower comparison chamber, the pure air filling the upper, and then measur-

ing the actual displacement for the varying concentrations.
The method is said to give an accuracy equivalent to .05%
of carbon dioxide.

METHOD OF
OBTAINING ALVEOLAR AIR.

ing the actual displacement for the varying concentrations.
The method is said to give an accuracy equivalent to .03%
of carbon dioxide.

METHODS OF OBTAINING ALVEOLAR AIR.

Two types of procedure are possible in the obtaining of samples of Alveolar air, giving, according to the procedure followed, air which is in approximate equilibrium with the partial Alveolar tension of the arterial blood in the pulmonary artery or air which is in approximate equilibrium with the partial Alveolar tension of the venous blood in the pulmonary capillaries. The former is obtained by a single forced expiration, the latter by repeated re-breathing of the air.

METHODS OF OBTAINING ALVEOLAR AIR.

Several methods for the first method are available. The following are the most commonly used. The first is the method of Haldane, of course, by this group. Haldane's is perhaps the simplest. This method uses a piece of glass tube with a glass mouth piece at one end. The subject breathes through the tube closing the mouthpiece with the tongue at the end of the expiration. The sample of air is drawn from the end of the tube. The other form uses valves for closing the tube, a necessary modification with operational subjects. Experiments carried out by Haldane show that a good sample of alveolar air can be obtained by this method. He observed, first, that the frequency of breathing was increased within a few minutes without altering the Alveolar tension of the gas percentage, provided that one depth of breathing is allowed to expire itself naturally, with a few depths of shallow breathing.

1. Haldane. J. Physiol.
2. A. Krogh and J. Lindhard. J. Physiol.

NO. 300724

RECEIVED AIR MAIL

METHODS OF OBTAINING ALVEOLAR AIR.

Two types of procedure are possible in the obtaining of samples of Alveolar air, giving, according to the procedure followed, air which is in approximate equilibrium with the carbon dioxide tension of the arterial blood in the pulmonary artery or air which is in approximate equilibrium with the carbon dioxide tension of the venous blood in the pulmonary capillaries. The former is obtained by a single forced expiration, the latter by repeated re-breathing of the air.

Several types of apparatus for the first method are available. Haldane (1) Lindhard and Krogh (2) have designed apparatus for the collection of single expirations. The Frid-ericia tube belongs, of course, to this group. Haldane's is perhaps the simplest. This method uses a piece of garden hose with a glass mouth piece at one end. The subject blows through the tube closing the mouthpiece with the tongue at the end of the expiration. The sample of air is drawn from the end of the tube. The other forms use valves for closing the tube, a necessary modification with untrained subjects. Experiments carried out by Haldane show that a true sample of Alveolar air can be obtained by this method. He observes, First: that the frequency of breathing may be varied within wide limits without altering the Alveolar Carbon Dioxide percentage, provided that the depth of breathing is allowed to regulate itself naturally, with no forcing or holding back.

- | | | |
|------------------------------|---------------|--------|
| 1. Haldane. | Am. Jr. Phys. | (1915) |
| 2. A. Krogh and J. Lindhard. | Jr. Phys. | (1914) |

METHODS OF OBTAINING ALVEOLAR AIR.

Two types of procedure are possible in the obtaining of samples of alveolar air, viz., according to the procedure followed, air which is in approximate equilibrium with the carbon dioxide tension of the arterial blood in the pulmonary artery or air which is in approximate equilibrium with the carbon dioxide tension of the venous blood in the pulmonary capillaries. The former is obtained by a single forced expiration, the latter by repeated re-breathing of the air.

Several types of apparatus for the first method are available. Haldane (1) and Krogh (2) have designed apparatus for the collection of single expirations. The first of these belongs, of course, to this group. Haldane's is perhaps the simplest. This method uses a piece of rubber hose with a glass mouth piece at one end. The subject blows through the tube closing the mouthpiece with the tongue at the end of the expiration. The sample of air is drawn from the end of the tube. The other form uses valves for closing the tube, a necessary modification with untimed subjects. Experiments carried out by Haldane show that a true sample of alveolar air can be obtained by this method. He observed, first, that the frequency of breathing may be varied within wide limits without affecting the alveolar carbon dioxide percentage, provided that the depth of breathing is allowed to remain itself naturally, with no forcing or

3/4 Respiration per minute.	Percentage of carbon dioxide.
30	5.67
4	5.66
36	5.68
3	5.71
24	5.48
6	5.56
4	5.72

Second: That when during normal breathing, air is expelled sharply from the lungs, the partial pressure of carbon dioxide in the expired air is constant after a certain volume has been expelled.

Depth of Expiration.	Percentage of Carbon dioxide.
190cc	3.03
335cc	4.37
510cc	5.04
650cc	5.19
950cc	5.51
1350cc	5.48

Also, he found that the mean of six successive determinations with a 900cc expiration showed a carbon dioxide percentage of 5.39 while a similar mean with an expiration of 1750cc showed a carbon dioxide percentage of 5.36. The deeper part of the expiration contains no more carbon dioxide than the middle part. Since the normal exchange of air is above 1000cc, it is evident that a fair sample of alveolar air is obtained by this method.

For the obtaining of samples of air by rebreathing, the Plesch method or one of its modifications is used. The apparatus and procedure used in Higgins' modification of the Plesch method is as follows: "To a copper disk of about 20cm. is soldered on the underside a flange 2cm wide. In the center of the same side of the disk is soldered an ordinary

Respiration per minute. Percentage of carbon dioxide.

30	5.57
4	5.55
25	5.55
5	5.71
25	5.48
5	5.55
4	5.75

Second: That when during normal breathing, air is expelled sharply from the lungs, the partial pressure of carbon dioxide in the expired air is constant after a certain volume has been expelled.

Table of Expiration. Percentage of Carbon dioxide.

150cc	5.55
250cc	5.57
350cc	5.55
450cc	5.55
550cc	5.55
650cc	5.55
750cc	5.55

Also, he found that the mean of six successive determinations with a 300cc expiration showed a carbon dioxide percentage of 5.55 while a similar mean with an expiration of 150cc showed a carbon dioxide percentage of 5.55. The deeper part of the expiration contains no more carbon dioxide than the middle part. Since the normal exchange of air is about 1000cc, it is evident that a fair sample of alveolar air is obtained by this method.

For the obtaining of samples of air by rebreathing,

the Fleisch method or one of its modifications is used. The apparatus and procedure used in Higgin's modification of the Fleisch method is as follows: "To a copper disk of about 80cm. is soldered on the underside a 1 1/2 inch 3/4 in wide. In the center of the same side of the disk is soldered an ordinary

3/4 inch three-way brass valve. Near this is soldered a small brass tap suitable for the attachment of small rubber tubing for taking samples of air. Over the top of the disk is stretched a rubber bathing cap, held in place by a large rubber band. The subject, with his nose clamped, breathes through a mouthpiece which is attached by a short piece of wide-bore rubber tubing to the three-way in the bottom of the copper disk. The tap is successively turned so that the subject first breathes to the outer air and then into the bag. After a definite amount of rebreathing, it is turned again, the bag is closed and a sample of the air is drawn from the bag for analysis." Different samples taken from the bag will show variations in the tension of carbon dioxide unless the air is thoroughly mixed by kneading the bag gently. More factors would seem to influence air obtained by this method than by that of Haldane. The tension of the carbon dioxide in the bag varies considerably with the depth of the initial inspiration and thus with the amount of air in the system of respiratory tract plus bag.

Moderately deep initial inspiration-

Carbon dioxide tension = 43.2 mm

Deeper initial inspiration-

Carbon dioxide tension = 41.5 mm

Very deep initial inspiration-

Carbon dioxide tension = 38.2 mm

Haldane (single expiration)

Carbon dioxide tension = 41.2 mm

This source of error can be to a degree avoided by adopting the suggestion of Higgins and filling the bag with a constant volume of air. The tap is turned at the end of a normal

expiration and the patient begins by inspiring from the bag. The time during which the breathing takes place is also a factor. After breathing from a bag four times in 20" the carbon dioxide tension was found to be 39.5 mm. After an interval of 3 1/2 minutes the same air was rebreathed for two periods of 20" each, the periods separated by an interval of one minute. The carbon dioxide tension then was 45.9 mm. After rebreathing for two more similar periods the carbon dioxide tension had risen to 46.4 mm. Continuation of the length of a single period will likewise cause a rise in the carbon dioxide tension.

4	respirations	in 20"	Carbon dioxide=	45.2 mm
6	"	" 30"	" "	= 47.4 mm
8	"	" 40"	" "	= 48.6 mm

The rate and depth of respiration while rebreathing is still another factor.

5	respiration	in 25"	Moderately deep	-Carbon dioxide=	46.4 mm
16	"	" "	Shallow	" "	=45.9 mm
5	"	" "	Very deep	" "	=46.3 mm
17	"	" "	Shallow	" "	=45.4 mm
5	"	" "	Natural	" "	=39.0 mm
5	"	" "	Moderately deep	" "	=45.1 mm

With the exception of the fifth determination in which the subject was breathing naturally, quietly and superficially, the values of the carbon dioxide tension agree well, whether the respirations were very deep or were rapid and shallow. Gentle superficial respiration does not suffice to bring about complete diffusion of gases. The same has been found to be true of very rapid superficial, panting breathing which also gives too low values for the carbon dioxide tension. Respirations of moderate depth seem to bring about as complete diffusion as very deep respirations. Best results have been obtained

expiration and the patient begins by inspiring from the bag. The time during which the breathing takes place is also a factor. After breathing from a bag four times in 30" the carbon dioxide tension was found to be 32.5 mm. After an interval of 3 1/2 minutes the same air was re-breathed for two periods of 30" each, the periods separated by an interval of one minute. The carbon dioxide tension then was 45.5 mm. After re-breathing for two more similar periods the carbon dioxide tension had risen to 46.4 mm. Continuing action of the lungs at a single period will likewise cause a rise in the carbon dioxide tension.

4	respiration in 30"	Carbon dioxide = 45.5 mm
3	"	" " = 47.1 mm
2	"	" " = 48.6 mm

The rate and depth of respiration while re-breathing is still another factor.

5	respiration in 30"	Moderately deep	Carbon dioxide = 45.4 mm
4	"	"	" " = 45.9 mm
3	"	Shallow	" " = 45.8 mm
2	"	Very deep	" " = 46.4 mm
1	"	Shallow	" " = 46.4 mm
0	"	Natural	" " = 39.9 mm
0	"	Moderately deep	" " = 45.1 mm

With the exception of the fifth determination in which the subject was breathing naturally, quietly and superficially, the values of the carbon dioxide tension agree well, whether the respirations were very deep or were rapid and shallow. Superficial respiration does not suffice to bring about complete diffusion of gases. The same has been found to be true of very rapid superficial, panting breathing which also gives too low values for the carbon dioxide tension. Respirations of moderate depth seem to bring about as complete diffusion as very deep respirations. Best results have been obtained

with five rather deep inspirations in twenty-five seconds. Series of observations which have compared results obtained by the Plesch method with those obtained by the method of Haldane on the same subject at the same time, show that the Plesch method gives results from one to four millimeters higher than the Haldane. It is claimed by some workers that the advantages of the Plesch-Higgins method are obvious. That since it is difficult to make sick people give an expiration of maximum depth and since anyone can breathe deeply enough to give accurate results with the rebreathing procedure, its use is to be preferred. According to Haldane's results, however, a breath of maximum depth is not required, and the author has found that the dislike of the patient for the rebreathing procedure, requiring as it does the closing of the nostrils, influences unfavorably the obtaining of the Alveolar air by that method.

The possibility of the diffusion of gases through rubber is potentially another source of error where the gas is collected in a thin walled rubber bag. The work of Dewar, however, would indicate that it could not be a disturbing factor where the time between collection and analysis was of brief duration.

The air for the Marriott determination is collected by the Plesch method. The rubber bag, however, as furnished by the dealers is an ordinary football bladder with a very narrow bore inlet tube. Experience has shown that such a narrow inlet prejudices seriously the efficacy of the equilibrium establishment and as a result we have substituted a modification of the Plesch Higgins apparatus just described.

with five rather deep inspirations in twenty-five seconds.
Series of observations which have compared results obtained
by the Fleisch method with those obtained by the method of
Haldane on the same subject at the same time, show that the
Fleisch method gives results from one to four millimeters
higher than the Haldane. It is claimed by some workers that
the advantages of the Fleisch-Higgins method are obvious.
That there is a difficulty in making such comparisons is
evident of maximum depth and since anyone can breathe deeply
enough to give accurate results with the Fleisch-Higgins
method, its use is to be preferred. According to Haldane's
results, however, a depth of maximum depth is not required,
and the author has found that the results of the patient
for the Fleisch-Higgins method, regarding as it does the clos-
ing of the nostrils, influences unfavorably the obtaining
of the alveolar air by that method.
The possibility of the diffusion of gases through
rubber is completely another source of error where the
gas is collected in a thin walled rubber bag. The work of
Cowan, however, would indicate that it could not be a dis-
turbing factor where the time between collection and analysis
was of brief duration.
The air for the Harriott determination is collected
by the Fleisch method. The rubber bag, however, as furnished
by the Harriott is an ordinary football bladder with a very
narrow bore inlet tube. Experience has shown that such a
narrow inlet prejudices seriously the efficacy of the equi-
librium establishment and as a result we have substituted a
modification of the Fleisch-Higgins apparatus just described.

A base for the cap is spun out of aluminum. The edge of this is rolled, serving to hold the cap in place without the use of the rubber band. The three-way brass valve is replaced by a protruding nipple an inch and a quarter in diameter. In this is inserted a rubber stopper thro which a short $3/4$ inch glass tube is passed. This tube is connected by thin walled rubber tubing to the mouth-piece, blown from glass tubing of the same bore. The bag is closed by a pinchcock on the rubber tubing. Samples of air are withdrawn thro this same outlet. In cases where patients are not able to cooperate, a mask covering the mouth and nose makes the collection of air possible.

C O N C L U S I O N .

C O N C L U S I O N .

The character of the study imposes certain conditions as criteria for the weighing of the methods to be employed. As has already been stated, a method to be available for the study, must offer, first: simplicity of operation; second: minimum of time consumption; and third; a sufficient degree of accuracy to warrant confidence in the generalizations to be based upon the measurements.

While none of these factors may be taken for granted and must all be subjected to experimental proof, the third, namely the accuracy of the method, takes the most important place. That just standards may be obtained for comparative purposes, it becomes necessary to adopt a standard method for the determination of Carbon dioxide, and one for the Ammonia in the Urine.

In regard to the first, the wealth of experimental evidence available coupled with a critical analysis of the technique of the method itself, leads to the selection of the Haldane apparatus as one which will give dependable and accurate results.

With the Urinary Ammonia on the other hand the field is by no means so limited, and several methods present themselves, all offering potential accuracy but involving equally possible sources of error. It may be assumed from the conditions of experiment that direct determination in the original solution, where the latter is of so highly complex a nature as is the Urine, offers so

large a number of possible disturbing factors as to render dependence upon it hardly warranted.

Of the indirect methods those involving the use of heat would again seem to be less desirable than those carried on at room temperature, owing to the possibly greater activities of the reagents upon other Urine constituents at the high temperature. By exclusion, one arrives at the Aeration and Precipitation methods and of ^{the} these latter, owing to the variability of the essential reagent, is, as has already been mentioned, susceptible to possibly grave error. Of the Aeration methods, the microchemical procedure of Folin-McCallum would seem to give most satisfactory results, both theoretically and practically. The speed of manipulation, and dependability of the same, all point to this method as being best suited to the purposes required.

Standards having been thus determined, the next point is selection of test methods which from their simplicity and rapidity seem to offer the greatest advantage. It is the purpose of the work in process to ascertain the relative accuracy of such methods and so beyond a brief examination of those sources of error, which are obviously incident to the procedure, the first two criteria can be assumed to be the determining factors in the choice.

Considering first the gas analysis, it is evident that a method in which but a single piece of apparatus is needed, offers decided advantage, other things being equal. As was previously stated, the Fridericia tube was designed

specifically for this purpose and naturally offers one of the methods to be investigated. In order to secure samples of gas from this apparatus for exact analysis by the Haldane apparatus, the instrument has been modified by the author in the following way: The large cock which is closed at the end of expiration thereby sealing the sample within the measuring tube, has been replaced by a three-way cock, the third arm being of capillary dimensions. After the collection of a sample within the tube and the equalization of the temperature of the gas by the usual procedure, mercury can be admitted at the bottom of the apparatus and a sample of the gas passed through the capillary tube into the Haldane apparatus and there subjected to exact analysis. The large bore of the ordinary delivery arm of the apparatus makes this modification necessary as the amount of air contained within it would prove a serious contaminator of the sample and vitiate the analysis made of the mixture. Of other methods available, the Gravimetric and Thermometric are quite obviously unsuitable. The volumetric method involves standard solutions, burettes, and other accessory pieces of apparatus which makes it not readily portable. Marriott's colorimetric method, on the other hand, offers decided advantage from its portability and simplicity of manipulation and sound theoretical basis. The method of collection of the air sample for this apparatus and naturally for comparison

with the Haldane instrument has already been discussed, the errors incident to the collection considered. The apparatus used by the author for this purpose is, as has already been stated, no more than a modification of the Higgins apparatus and possesses the only advantage over the latter of lightness and slightly simpler manipulation. It is to be regretted that comparative measurements could not be made with the Interferometer, but the instrument has not been available up to the present time, although it is hoped in some of the later series to make use of it. The accuracy calculated, namely .05%, is equal to about 1% of the total amount, and which, as the measurements are reported in terms of millimetres of mercury pressure and vary from 15 to 45, would assume a wholly negligible magnitude.

The work then comprises the determination of the Alveolar Carbon dioxide by means of the Fridericia apparatus, with control determination by the Haldane, on the tension of the gas in arterial blood, and determination on the same patient of the venous tension by means of the Mariott method with the Haldane again being used for control, the various factors of importance as already outlined determined by experiment.

The selection of a method for the determination of Urinary Ammonia is a matter of some difficulty. The direct Nesslerization of the Urine as suggested by Folin and Denis, and Sumner, the direct titration or the so-called "formol" method, and the precipitation method suggested by

Folin and Bell, all present points of desirability. As, however, the two methods suggested by Folin require the use of a colorimeter, an instrument which even in its less expensive forms calls for a considerable outlay of money, ^{this} renders them the less desirable from the standpoint of the clinician. The Ronchèse-Malfatti or "formol" titration while, as has been pointed out, ~~is~~ susceptible of certain inherent errors, does not call for an elaborate apparatus and is both rapid and simple. The tenth normal caustic soda which is required is also used for certain other laboratory procedures incident to clinical diagnosis and hence forms a part of even modest laboratory equipment. For these various reasons it was selected as the single comparative method for Urine Ammonia.

The character of the Folin-McCallum aeration is such as to bring it also within the possible range of clinical procedure. This phase of study then consists of the determination of Ammonia content in the 24 hour elimination. As the total amount of the Ammonia is to be regarded as less of an index than the amount in relation to the total nitrogen, this latter factor has also been determined as a matter of routine adopting the thoroughly familiar Kjeldahl method. A sample of the form used for report is here included, and shows the scope of the experimental work as it is being carried out.

In conclusion, the writer wishes to say that in presenting this dissertation, no attempt has been made to include experimental results, as these are at the present

moment in process of compilation and have not reached a point in progress to warrant any conclusions. It is purposed at some later date to present these experimental results and the conclusions drawn from them.

RECEIVED

moment in process of completion and have not reached a
point in progress to warrant any conclusions. It is
proposed at some later date to present these experimental
results and the conclusions drawn from them.

BIBLIOGRAPHY

James Ewing and J. H. H. H. H.
The Effect of the Metabolism of Fatty Acids
on the Metabolism of the Fetus in Pregnancy.
American Journal of Obstetrics 25 255 (1923)

S. G. van Vliet
"Metabolism of Amino Acids in Pregnancy".
Biochimica et Biophysica Acta 11 181-85 (1953)

P. J. Underhill and E. H. H.
"Metabolism of Fatty Acids in Pregnancy".
Archives of Internal Medicine 5 25-31 (1924)

Dorothy E. Lindsay
"The Distribution of Metabolism in the Fetus".
Biology and the Fetus (1924) 1-10
BIOBIOGRAPHY . 12

E. C. Bailey
"Protein Metabolism in Late Pregnancy".
Journal of the American Medical Association 11 181-85 (1923)

Th. Heynemann
"The Question of the Liver and Crystalline Metabolism
during Pregnancy and in the Fetus of Pregnancy".
Zeitschrift für Biologie und Anatomie 13 217 (1923)

H. Schirckner
"Carbohydrate Metabolism during Pregnancy".
Berliner klinische Wochenschrift 22 202-3 (1923)

J. Smith
"Carbohydrate Metabolism during Pregnancy and in
the Fetus".
Monatsschrift für Geburtshilfe und Gynäkologie 57 413 (1923)

J. R. Hurler and E. C. Bailey
"Further Observations on Protein Metabolism of
Normal Pregnancy".
Archives of Internal Medicine 12 25-31 (1924)

E. A. Hasselbach and E. A. Gammeltoft
"Metabolism of Fatty Acids in Pregnancy".
Biochimica et Biophysica Acta 11 181-85 (1953)

BIBLIOGRAPHY.

PREGNANCY.

METABOLISM

- James Ewing and C.G.L.Wolf
"The Clinical Significance of Urinary Nitrogen.
II The Metabolism in the Toxemias of Pregnancy".
American Journal of Obstetrics 55 289 (1907)
- E.C.van Tiersen
"Excretion of Amino Acids in Pregnancy".
Biochemie Zeitschrift 11 121-36 (1909)
- F.P.Underhill and R.F.Rand
"Peculiarities of Nitrogenous Metabolism in Pernicious Vomiting of Pregnancy".
Archives of Internal Medicine 5 61-91 (1910)
- Dorothy E. Lindsay
"The Distribution of Nitrogen in the Maternal Urine and in the Fetal Fluids throughout Pregnancy".
Biochemical Journal 6-76 (1912)
- H.C.Bailey
"Protein Metabolism in Late Pregnancy".
Journal of the American Medical Association 59 1522 (1913)
- Th. Heynemann
"The Question of the Liver and Creatinine Metabolism during Pregnancy and in the Toxicoses of Pregnancy".
Zentralblatt für Biochemie und Biophysik 13 817 (1913)
- H.Schirokaner
"Carbohydrate Metabolism during Pregnancy".
Berliner klinische Wochenschrift 49 500-3 (1913)
- J. Beuthin
"Carbohydrate Metabolism during Gravidity and in Eclampsia".
Monatschrift für Geburtshilfe und Gynakologie 37 413 (1913)
- J.R.Murlin and H.C.Bailey
"Further Observations on Protein Metabolism of Normal Pregnancy".
Archives of Internal Medicine 12 288-314 (1913)
- K.A.Hasselbalch and S.A.Gemmeltøft
"Neutrality Regulation of the Pregnant Organism".
Biochemie Zeitschrift 68 206-64 (1915)

METABOLISM (continued)

L.A.Emge

- "Acidosis in Normal Uterine Pregnancy".
American Journal of Obstetrics 74 769 (1918)

TOXEMIAS

J.W.Williams

- "Pernicious Vomiting of Pregnancy".
Bulletin of Johns Hopkins Hospital 17 71 (1906)

James Ewing and C.G.L.Wolf

- "The Metabolism in the Toxemia of Pregnancy".
American Journal of Medical Science 55 289 (1907)

F.P.Underhill and R.F.Rand

- "Peculiarities of Nitrogenous Metabolism in
Pernicious Vomiting of Pregnancy".
Archives of Internal Medicine 5 61-91 (1910)

James Ewing

- "Toxemia of Pregnancy".
American Journal of Medical Science 139 828(1910)

G.C.Trawick

- "Toxemias of Pregnancy".
Southern Medical Journal 3 416-20 (1910)

W.L.Crosthwait

- "The Toxemias of Pregnancy with Special
Reference to the Pre-eclamptic State".
Texas State Journal of Medical Science
5 341 (1909-10)

Ross McPherson

- "A Consideration of the Convulsive Toxemias
of Pregnancy".
International Journal of Surgery 23 116-119 (1910)

E. Landsberg

- "Urine and Blood in Eclampsia".
Gynakologie und Geburtshilfe 73 234-66 (1913)

H. Williamson

- "Pregnancy Toxemia - A Study of Acidosis
in Pregnancy".
American Medicine 11 385-92 (1916)

J.R.Losee and D.D. Van Slyke

- "Acidosis in Toxemias of Pregnancy".
Journal of American Medical Association
68 659 (1917)

"The Toxemias of Pregnancy".

- American Journal of Medical Science.
94-100 (1917)

TOXEMIAS (continued)

- Frank W. Lynch
"Treatment of the Severe Vomitings of Early
Pregnancy".
Journal of the American Medical Association 73 488 (1919).
- Clifford Mitchell
"Toxemias of Pregnancy".
Hahnemannian Monthly 52 641-8 (1918)
- Isei Obata
"The Nature of Eclampsia".
Journal of Immunology 4 111 (1919)
- J.W. Williams
"Toxemias of Pregnancy".
Chapter 26 "Obstetrics" 4th Edition (1917)
Published by Appleton and Co. New York and London
- H.G. Wells
"Toxemias of Pregnancy".
"Chemical Pathology" 533 3rd Edition (1918)
Published by W.B. Saunders and Co. Philadelphia and
London.
- J.M. Slemans
"Analysis of the Blood in Eclampsia and Allied
Autointoxications".
Journal of the American Medical Association 70 1793 (1918)

RESPIRATION.

- K.A. Hasselbach
"The Physiology of Respiration during Pregnancy".
Scandinavisches Archiv für Physiologie 27 12 (1913)
- A. Leimdorfer, J. Novak and O. Porges
"Carbon Dioxide Tension of the Blood in Pregnancy".
Zeitschrift für klinische Medizin 80 13 (1913)
- S.H. Blodgett
"Laboratory Aids to the Obstetrician".
Journal of the American Institute of Homeopathy. 108 94-8 (1918)

AMMONIA COEFFICIENT.

- A.W. Ellis
"The Ammonia Coefficient in a case of Severe
Vomiting in Pregnancy".
Journal of the Canadian Medical Association 59 1522 (1913)

AMMONIA COEFFICIENT (continued)

- A.A.Hotaling
"The Ammonia Coefficient in Pregnancy".
American Journal of Obstetrics 67 925-30 (1913)
- S.H.Blodgett
"Laboratory Aids to the Obstetrician".
Journal of the American Institute of Homeopathy
108 94 (1918)
- J.R.Garber
"Significance of the Ammonia Coefficient in
Obstetrical Work".
Southern Medical Journal 9 900 (1918)

ACIDOSIS.

- O. Folin
"The Acid Intoxication Theory".
Transactions of the Association of American
Physicians. 22 256 (1907)
- E.P.Joslin
"Acidosis".
Boston Medical and Surgical Journal 157 40-43(1907)
- James Ewing
"Acidosis and Associated Conditions".
Archives of Internal Medicine 2 330-54 (1909)
- "A Classification of Acidosis".
Transactions of Association of American Physicians
23 31-8 (1909)
- A.W.Sellards
"Essential Features of Acidosis and their occurrence
in Chronic Renal Disease".
Bulletin of Johns Hopkins Hospital 25 141 (1914)
- Marius Lauritzen
"The Determination of Acidosis and Its Clinical
Application in Diabetes Mellitus".
Zeitschrift fur klinische Medizin 80 13 (1914)
- Yandell Henderson
"The Time That the Breach Can Be Held as an Index
of Acidosis".
Journal of the American Medical Association
63 318 (1914)
- F.W.Peabody
"Studies on Acidosis and Dyspnea in Renal and
Cardiac Diseases".
American Journal of Medical Science 14 236-262
(1914)

ACIDOSIS (continued)

- H.Steenbock, V.E.Nelson and E.B.Hart
 "Acidosis in Omnivora and Herbivora and Its
 Relation to Protein Storage".
 Journal of Biological Chemistry 19 399-419 (1915)
- A.P.Beddard, M.S.Pembrey and E.I.Spriggs
 "Observations upon Acidosis in Diabetes Mellitus".
 British Medical Journal II 389 (1915)
- P.J.Cammidge
 "Diabetic Acidosis".
 American Medicine 11 363-73 (1916)
- J.Howland and W.Mck. Marriott
 "A Discussion of Acidosis".
 Johns Hopkins Hospital Bulletin 27 63-9 (1916)
- H.L.Higgins, F.W.Peabody and R.Fitz
 "The Study of Acidosis in Three Normal Subjects".
 Journal of Medical Research 34 263-72 (1916)
- L.G.Rowntree
 "The Diagnosis and Treatment of Acidosis".
 Transactions of the Association of American
 Physicians. 31 39-57 (1916)
- C.Frothingham, Jr.
 "Acidosis in Acute and Chronic Diseases".
 Archives of Internal Medicine 18 717-32 (1917)
- D.D.Van Slyke and E.Stillman and G.E.Cullen
 "The Nature and Detection of Diabetic Acidosis".
 Proceedings of the Society of Experimental
 Biology and Medicine 11 165-7 (1917)
- E.Stillman, D.D.Van Slyke, G.E.Cullen and R.Fitz
 "The Blood, Urine and Alveolar Air in Diabetic
 Acidosis".
 Journal of Biological Chemistry 30 405-56 (1917)
- J.P.Peters, Jr.
 "Carbon dioxide Acidosis the Cause of Cardiac
 Dyspnea".
 American Journal of Physiology 43 113-20 (1917)
- J.H.Austin and Leon Jonas
 "Clinical Studies of Acidosis".
 American Journal of Medical Sciences 153 81-94 (1917)
- A.W.Sellards
 "The Principles of Acidosis and Clinical Methods
 for Its Study".
 Published Harvard University Press, Cambridge (1917)

ACIDOSIS (continued)

- H. Steenbock, V. E. Nelson and E. E. Hart
"Acidosis in Canine and Ruminants and Its
Relation to Protein Metabolism".
Journal of Biological Chemistry 19 399-419 (1915)
- A. E. Bennett, M. S. Pomeroy and E. L. Spriggs
"Observations upon Acidosis in Diabetic Mellitus".
British Medical Journal II 389 (1915)
- E. J. Carr-Saunders
"Diabetic Acidosis".
American Medicine 11 353-73 (1915)
- J. Howard and W. McK. Harriott
"A Discussion of Acidosis".
Jones Hopkins Hospital Bulletin 27 63-9 (1915)
- H. I. Higgins, E. W. Peabody and R. Pitts
"The Study of Acidosis in Three Normal Subjects".
Journal of Medical Research 34 353-73 (1915)
- I. G. Sawastee
"The Metabolism and Treatment of Acidosis".
Transactions of the Association of American
Physicians 31 33-37 (1915)
- C. F. Probst, Jr.
"Acidosis in Acute and Chronic Diseases".
Archives of Internal Medicine 18 717-32 (1917)
- C. D. Van Slyke and E. A. Millman and C. E. Gillen
"The Nature and Detection of Diabetic Acidosis".
Proceedings of the Society of Experimental
Biology and Medicine 11 155-7 (1917)
- E. A. Millman, C. D. Van Slyke, C. E. Gillen and R. Pitts
"The Blood, Urine and Alveolar Air in Diabetic
Acidosis".
Journal of Biological Chemistry 30 405-36 (1917)
- J. F. Peters, Jr.
"Carbon Dioxide Acidosis the Cause of Cardiac
Dyspnea".
American Journal of Physiology 43 113-20 (1917)
- J. H. Austin and Leon Jones
"Clinical Studies of Acidosis".
American Journal of Medical Sciences 153 81-94 (1917)
- A. F. Bellis
"The Principles of Acidosis and Clinical Methods
for Its Study".
Published Harvard University Press, Cambridge (1917)

ACIDOSIS (continued)

- R.P.Gradwohl and A.J.Blaivais
 "Blood and Urine Chemistry" 173
 Published C.V.Mosby Co. St. Louis. (1917)
- D.D.Van Slyke
 "Studies of Acidosis".
 Journal of Biological Chemistry 33 271 (1918)
- V.W.Jensen
 "Acidosis in Relation to Pancreatic Diabetes".
 Journal of Biological Chemistry 33 8 (1918)
- J.R.MacLeod
 "The Spontaneous Development of an Acidosis Con-
 dition in Decerebrated Cats".
 Proceedings of the Society of Experimental and
 Biological Medicine. 16 28 (1918)
- H.L.McNeil and M.D.Levy
 "Blood and Urine Ammonia in Acidosis".
 Journal of Laboratory and Clinical Medicine (1919)
- J.J.R.MacLeod
 "Diagnosis of Acidosis".
 Journal of Laboratory and Clinical Medicine
 4 315 (1919)

ACID BASE EQUILIBRIUM.

- Andrew W. Sellards
 "The Determination of the Equilibrium in the
 Human Body between Acids and Bases with
 Especial Reference to Acidosis and Nephro-
 pathies".
 Johns Hopkins Hospital Bulletin 23 289 (1912)
- L.H.Newburgh, W.W.Palmer and L.J.Henderson
 "Clinical Studies on Acid Base Equilibrium
 and the Nature of Acidosis."
 Archives of Internal Medicine 12 153-70 (1913)
- J.Christiansen, C.G.Douglas, J.S.Haldane
 "The Absorption and Disassociation of Carbon
 Dioxide by Human Blood."
 Journal of Physiology 48 244 (1914)
- A.W.Sellards
 "A Clinical Method for Studying Titratable
 Alkalinity of the Blood and Its Application
 to Acidosis".
 Bulletin of Johns Hopkins Hospital 25 101 (1915)

ACIDOSIS (continued)

- (1917) H.F. Grubb and A.L. Bickel
"Blood and Urine Chemistry" 173
Published C.V. Mosby Co. St. Louis.
- (1918) O.D. Van Slyke
"Studies of Acidosis".
Journal of Biological Chemistry 33 271
- (1918) V.F. Jansen
"Acidosis in Relation to Pancreatic Diabetes".
Journal of Biological Chemistry 33 8
- (1918) J.R. MacLeod
"The Spontaneous Development of an Acidosis Con-
dition in Decapitated Rats".
Proceedings of the Society of Experimental and
Biological Medicine 15 28
- (1919) H.L. McNeill and M.C. Levy
"Blood and Urine Chemistry in Acidosis".
Journal of Laboratory and Clinical Medicine (1919)
- (1919) J.R. MacLeod
"Diagnosis of Acidosis".
Journal of Laboratory and Clinical Medicine 4 215

ACID BASE EQUILIBRIUM.

- (1913) Andrew W. Salter
"The Determination of the Equilibrium in the
Human Body between Acid and Base with
Special Reference to Acidosis and Nephro-
pathies".
Johns Hopkins Hospital Bulletin 23 289
- (1913) I.R. Newburgh, W.E. Palmer and L.F. Henderson
"Clinical Studies on Acid Base Equilibrium
and the Nature of Acidosis".
Archives of Internal Medicine 13 153-70
- (1914) J. Christensen, C.E. Dogliani, J.E. Williams
"The Absorption and Dissociation of Carbon
Dioxide by Human Blood".
Journal of Physiology 48 244
- (1915) A.W. Salter
"A Clinical Method for Studying Titratable
Acidity of the Blood and Its Application
to Acidosis".
Bulletin of Johns Hopkins Hospital 25 101

ACID BASE EQUILIBRIUM (continued)

K.A.Hasselbalch

"Reduced and Regulated Hydrogen Number of the Blood".

Biochemie Zeitschrift 74 56-62 (1916)

R.Fitz and D.D.Van Slyke

"The Relation between Alkaline Reserve and Acid Excretion".

Journal of Biological Chemistry 30 289 (1917)

J.F.McClendon, A.Shedlov and W.Thomson

"Tables for finding the Alkaline Reserve of Blood Serum".

Journal of Biological Chemistry 31 519 (1917)

D.D.Van Slyke and G.E.Cullen

"Bicarbonate Concentration of the Blood Plasma, Its Significance, and Its Determination as a Measure of Acidosis".

Journal of Biological Chemistry 30 289 (1917)

AMMONIA CONTENT OF THE URINE.

S.A.Gammeltoft

"The Ammonia Content of the Urine and its Relation to Gastric Secretion".

Zeitschrift für Physiologische Chemie 75 57 (1912)

Alonzo E. Taylor

"Output of Ammonia in Normal Urine".
Proceedings of the American Society of
Biological Chemistry

(1910)

L.Lematte

"Nitrogenous Metabolism".

Bulletin des sciences Pharmacolog 20 578 (1914)

K.A.Hasselbalch

"Ammonia as Physiological Neutrality Regulator."

Biochemie Zeitschrift 74 18 (1916)

W.Denis and A.S.Minot

"Ammonia Excretion as Influenced by the Ingestion of Alkali".

Journal of Biological Chemistry 35 101 (1918)

H.C.McNeil and M.D.Levy

"Observations on the Blood and Urine Ammonia in Acidosis".

Journal of Laboratory and Clinical Medicine 3 18 (1918)

ACID-BASE EQUILIBRIUM (continued)

- Y. A. Hasegawa
"Effect of Regulated Hydrogen Ion Concentration on the Blood"
Biochimica et Biophysica Acta 55-56 74 (1973)
- E. Witz and G. Van Slyke
"The Relation Between Alkaline Reserve and Acid Equivalents"
Journal of Biological Chemistry 20 289 (1917)
- L. F. Hollander, A. Hedberg and W. Thorsen
"Tables for Finding the Alkaline Reserve of Blood Serum"
Journal of Biological Chemistry 21 219 (1917)
- G. C. Van Slyke and L. F. Hollander
"Alkaline Reserve of the Blood Plasma, Its Significance, and Its Determination as a Measure of Acidosis"
Journal of Biological Chemistry 20 289 (1917)

AMMONIA CONTENT OF THE URINE

- S. A. Gerasimov
"The Ammonia Content of the Urine and its Relation to Gastric Secretion"
Biochimica et Biophysica Acta 75 67 (1973)
- Alonso E. Taylor
"Output of Ammonia in Normal Urine"
Proceedings of the American Society of Biological Chemistry (1913)
- I. Jansz
"Ammonia Secretion"
Biochimica et Biophysica Acta 20 273 (1973)
- K. A. Hasegawa
"Ammonia as Physiological Metabolic Regulator"
Biochimica et Biophysica Acta 74 18 (1973)
- W. Denis and S. S. Klotz
"Ammonia Secretion as Influenced by the Location of Alkali"
Journal of Biological Chemistry 25 101 (1913)
- H. C. McNeill and M. C. Levy
"Observations on the Blood and Urine Ammonia in Acidosis"
Journal of Laboratory and Clinical Medicine 3 18 (1913)

URINARY AMMONIA. Methods of Determination.

A.Ronchese

"A New Method for the Determination of Ammonia".
Compte rendus Societe de Biologie 62 799 (1907)

H. Malfatti

"Clinical Methods for the Estimation of Ammonia".
Zeitschrift für analytische Chemie.47 273 (1908)

P.E.Howe and P.B.Hawk

"Comparative Tests of Spiro's and Folin's Methods
for the Determination of Ammonia".
Journal of Biological Chemistry 5 477 (1909)

R.F.Bacon

"A Rapid Clinical Method for the Determination
of the Ammonia Coefficient of Urines".
Phillipine Journal of Science 4A 153 (1909)

A.Florence

"Exact Gasometric Estimation of Urea and Ammonia".
Compte rendus Societe de Biologie 148 943 (1910)

G.Piccinnine

"Metabolism and Determination of Ammonia".
Riviste critica di clinica medica 11 5 (1910)

C.V.Spindler

"Titration of Ammonia by the Formol Method".
Schweiz Wochenschrift 47 767 (1909)

J.C.Tomlinson

"The Estimation of Free Ammonia and Ammonium
Carbonate".
Chemical News 103 13 (1911)

H.Bjorn Andersen and Marius Lauritzen

"Acid and Ammonia Estimation in Urine and their
Clinical Application".
Zeitschrift für Physiologie und Chemie 64 21 (1911)

O.Folin and C.E.Farmer

"A New Method for the Determination of Total Nitrogen
in the Urine".
Journal of Biological Chemistry 11 493 (1912)

L.Lematte

"Separation and Volumetric Determination of Urinary
Urea and Ammonia".
Compte rendus Societe de Biologie 174 217 (1913)

L.W.Winkler

"The Volumetric Determination of Ammonia".
Zeitschrift für angewandte Chemie 26 231 (1913)

UNIVERSITY MEMORIAL. Methods of Determination.

- A. Rönneberg
"A New Method for the Determination of Ammonia".
Compte Rendu Société de Biologie 62 729 (1907)
- H. Kallstedt
"Clinical Methods for the Estimation of Ammonia".
Zeitschrift für analytische Chemie 47 878 (1908)
- F. E. Howe and F. W. Hawk
"Comparative Tests of Spiro's and Folin's Methods
for the Determination of Ammonia".
Journal of Biological Chemistry 5 477 (1909)
- F. E. Hawk
"A Rapid Clinical Method for the Determination
of the Ammonia Content of Urine".
Philippine Journal of Science 44 125 (1909)
- A. Rönneberg
"Exact Gravimetric Estimation of Free and Ammonia".
Compte Rendu Société de Biologie 148 217 (1910)
- G. Nicolai
"Metabolic and Determination of Ammonia".
Ziveste original in clinical medicine 11 8 (1910)
- G. V. Seidner
"Estimation of Ammonia by the Folin Method".
Zeitschrift für analytische Chemie 47 767 (1909)
- L. C. Tomlinson
"The Estimation of Free Ammonia and Ammonium
Carbonate".
Chemical News 103 18 (1911)
- H. Bjorn Andersen and Martin Lauritzen
"Acid and Ammonia Estimation in Urine and their
Clinical Application".
Zeitschrift für Physiologie und Chemie 64 81 (1911)
- G. Folin and W. E. Warner
"A New Method for the Determination of Total Nitrogen
in the Urine".
Journal of Biological Chemistry 11 493 (1912)
- L. Rönneberg
"Separation and Volumetric Determination of Urinary
Urea and Ammonia".
Compte Rendu Société de Biologie 174 217 (1913)
- L. V. Finkler
"The Volumetric Determination of Ammonia".
Zeitschrift für analytische Chemie 50 231 (1913)

METHODS OF AMMONIA DETERMINATION (continued)

- R. Rosenbloom
"Clinical Methods for the Estimation of Total Nitrogen and Ammonia in the Urine".
Journal of the American Medical Association 61 87 (1913)
- P.A.Kober and S.S.Graves
"Quantitative Distillation by Aeration, for Kjeldahl, Urea and other Nitrogen Estimates."
Journal of the American Chemical Society 35 1594 (1913)
- A.Schulze
"Winkler's Volumetric Determination of Ammonia".
Mitteilunge konigliches Landesanstalt für Wasserhygiene 18 87 (1914)
- Koloman Budai
"Quantitative Determination of Ammonia and Trimethylamine".
Zeitschrift für Physiologische Chemie 86 107 (1914)
- Addison Gulick
"A Simplification of the Determination of Total Nitrogen by Colorimetry".
Journal of Biological Chemistry 18 541 (1914)
- L.W.Winkler
"Determination of Ammonia by the Boric Acid Method".
Zeitschrift für angewandte Chemie 271 630 (1915)
- S.S.Graves
"A Precipitant for Ammonia".
Journal of the American Chemical Society 37 1171 (1915)
- A.A.Bonnema
"A Simple Clinical Method for the Determination of Ammonia in the Urine".
Chemical Zeitung 39 519 (1915)
- J.C.Bock and S.R.Benedict
"An Examination of the Folin-Farmer Method for the Colorimetric Estimation of Ammonia".
Journal of Biological Chemistry 20 47 (1915)
- W.C.DeGraff
"Ammonia and Amino Acid in the Urine".
Pharm. Weekblad 52 1777-8 (1916)
- M.H.Scheltema
"The Determination of Ammonia, Amino Acids and Albumen in Urine by means of Formaldehyde".

METHODS OF AMMONIA DETERMINATION (continued)

H. Rosenblum
"Clinical Methods for the Determination of Total
Nitrogen and Ammonia in the Urine".
Journal of the American Medical Association

(1913) 87 51

E.A. Kober and E.E. Graves
"Quantitative Distillation by Aeration, for
Urethral, Urea and other Nitrogen Compounds".
Journal of the American Chemical Society

(1913) 35 1334

A. Schmitz
"Winkler's Volumetric Determination of Ammonia".
Mittelingen Koniglichen Laboratorium fur
Vergewisserte

(1914) 18 67

Kolomon Budai
"Quantitative Determination of Ammonia and
Trimethylamine".
Zeitschrift fur Physiologische Chemie

(1914) 96 197

Adrian Glick
"A Simplification of the Determination of Total
Nitrogen by Colorimetry".
Journal of Biological Chemistry

(1914) 18 241

E.W. Winkler
"Determination of Ammonia by the Boric Acid Method".
Zeitschrift fur angewandte Chemie

(1915) 27 633

E.E. Graves
"A Precipitant for Ammonia".
Journal of the American Chemical Society

(1915) 37 1171

A.A. Johnson
"A Simple Clinical Method for the Determination
of Ammonia in the Urine".
Chemical Abstracts

(1915) 39 218

J.C. Beck and E.E. Graves
"An Examination of the Folin-Carrier Method for the
Colorimetric Estimation of Ammonia".
Journal of Biological Chemistry

(1915) 30 47

F.C. Garrett
"Ammonia and Amino Acids in the Urine".
Bull. National

(1915) 32 1777-8

M.E. Schellberg
"The Determination of Ammonia, Amino Acids and
Alcohols in Urine by means of Formaldehyde".

METHODS OF AMMONIA DETERMINATION (continued)

- O.Folin and W.Denis
"Ammonia in Urine".
Journal of Biological Chemistry 26 497 (1916)
- E.Bernard
"Titrimetric Determination of Ammonia According
to L.W.Winkler".
Zentralblatt für Biochemie und Biophysik (1916)
- G.E.Foxwell
"A New Method for Estimating Ammonia".
Gas World 64 1654 (1916)
- P.A.Kober
"Efficiency of the Aeration method for Distilling
Ammonia".
Journal of the American Chemical Society 38 2568 (1916)
- Ch. Schenitzky
"Ammonia Determination in Human Urine".
Biochemie Zeitschrift. 76 177 (1916)
- W.C.deGraff and J.E.Van der Zande
"Ammonia and Amino Acids in the Urine".
American Journal of Physiology 6 230 (1903)
- O.Folin and R.D.Bell
"Application of a new Reagent for the Separation
of Ammonia".
Journal of Biological Chemistry 29 329 (1917)
- R.Weiss
"Quantitative Determination of Ammonia and Sulfur".
Munchener Medizinische Wochenschrift 63 1547 (1918)
- A.Leclere
"Determination of Urinary Ammonia".
Compte rendus Societe de Biologie 80 959 (1918)
- J.B.Sumner
"A New Method for the Direct Nesslerization
of Ammonia in the Urine".
Journal of Biological Chemistry 34 37 (1918)
- Neubauer
Journal of Praktische Chemie 64 278 (1855)
- J.Kjeldahl
"New Method for the Determination of Nitrogen
in Organic Substances".
Zeitschrift für Analytische Chemie 23 366 (1883)

METHODS OF AMMONIA DETERMINATION (continued)

- G. Folin and W. Denis
"Ammonia in Urine".
Journal of Biological Chemistry 33 437 (1916)
- E. Bernhardt
"Titrimetric Determination of Ammonia according to E. W. Folin".
Centralblatt für Bakteriologie und Supplement (1916)
- G. E. Foxwell
"A New Method for Estimating Ammonia".
Gas World 54 1554 (1916)
- F. A. Mohr
"Efficiency of the Kjeldahl method for Distilling Ammonia".
Journal of the American Chemical Society 38 1863 (1916)
- G. A. Gohenschalk
"Ammonia Determination in Human Urine".
Biochemische Zeitschrift 75 177 (1916)
- W. G. Bostick and E. M. Van der Bunde
"Ammonia and Amino Acids in the Urine".
Biochemische Zeitschrift 75 177 (1916)
- G. Folin and W. G. Bell
"Application of a new Reagent for the separation of ammonia".
Journal of Biological Chemistry 33 323 (1917)
- E. Weiss
"Quantitative Determination of Ammonia and Sulphur".
Mannheimer Medizinische Wochenschrift 53 1547 (1918)
- A. Jochims
"Determination of Urinary Ammonia".
Zeitschrift für klinische Chemie 80 932 (1918)
- J. B. Sumner
"A New Method for the Direct Nesslerization of Ammonia in the Urine".
Journal of Biological Chemistry 34 37 (1919)
- H. H. Sumner
"Journal of Praktische Chemie 34 278 (1922)
- J. Kjeldahl
"New Method for the Determination of Nitrogen in Organic Substances".
Zeitschrift für Analytische Chemie 33 366 (1883)

METHODS OF AMMONIA DETERMINATION (continued)

- Neubauer und Huppert
 "Analyse des Harns"
 Published C.W.Kreidels, Weisbaden (1910)
- Schlösing
 "The Determination of Ammonia".
 Annales de Chimie et de Physique 31 153 (1851)
- Boussingault
 "Determination of Ammonia".
 Annales de Chimie et de Physique 29 472 (1850)
- C.Würster
 "Ammonia determination in Urine".
 Centralblatt für Physiologie 1 485 (1887)
- Steyer
 "Determination of Ammonia".
 Zeit. zur Chem. Phys. und Path. 11 312 (1902)
- Kruger, Reich and Schittenhelm
 "Determination of Ammonia".
 Zeitschrift für Physiologische Chemie 39 165 (1903)
- Phillip Schaffer
 "Quantitative Determination of Ammonia in Urine."
 American Journal of Physiology 8 330 (1903)
- O.Folin
 "A New Method for the Determination of Ammonia in
 Urine and Other Animal Fluids".
 Zeitschrift für Physiologische Chemie (1902)
- Steele
 "Modification of the Folin Method in the Presence
 of Magnesium".
 Proceedings of the Society of Experimental Biology
 6 127 (1909)
- Steele and Gies
 "Notes of Effects of Folin Method in the Presence
 of an Excess of Magnesium in the Urine".
 Journal of Biological Chemistry 5 71 (1909)
- S.R.Benedict and Emil Osterberg
 "A Note on the Determination of Ammonia in the Urine".
 Biochemical Bulletin 3 41 (1913)
- O.Folin
 "Note of the Determination of Ammonia in the Urine".
 Journal of Biological Chemistry 8 497 (1910)
- R.O.Davis
 "The Determination of Ammonia without a Condensor".
 Journal of the American Chemical Society
 31 556 (1909)

METHODS OF AMMONIA DETERMINATION (continued)

- Neubauer and Hubert
"Analyse des Harns"
Verlag J. F. Schöner, Leipzig
(1910)
- Schilling
"The Determination of Ammonia"
Annales de Chimie et de Physique 31 153
(1911)
- Bosselagault
"Determination of Ammonia"
Annales de Chimie et de Physique 33 432
(1912)
- C. W. Hunter
"Ammonia Determination in Urine"
Centralblatt für Bakteriologie 1 435
(1913)
- Bayer
"Determination of Ammonia"
Zeit. für Chem. Phys. und Path. 11 312
(1913)
- Kirner, Jochen and Schilling
"Determination of Ammonia"
Centralblatt für Bakteriologie 33 155
(1913)
- Philip Schiller
"Quantitative Determination of Ammonia in Urine"
American Journal of Physiology 8 380
(1913)
- O. Polin
"A New Method for the Determination of Ammonia in Urine and Other Animal Fluids"
Centralblatt für Bakteriologie Chemie 33 155
(1913)
- Steele
"Modification of the Polin Method in the Presence of Magnesium"
Proceedings of the Society of Experimental Biology 6 137
(1913)
- Steele and Glas
"Notes on Effects of Polin Method in the Presence of an Excess of Magnesium in the Urine"
Journal of Biological Chemistry 3 41
(1913)
- E. J. Benedict and Emil Osterberg
"A Note on the Determination of Ammonia in the Urine"
Biochemical Bulletin 3 41
(1913)
- O. Polin
"Note of the Determination of Ammonia in the Urine"
Journal of Biological Chemistry 3 437
(1913)
- E. O. Davis
"The Determination of Ammonia without a Condenser"
Journal of the American Chemical Society 31 355
(1909)

METHODS OF AMMONIA DETERMINATION (continued)

- O.Folin and A.B.MaCallum
 "The Determination of Ammonia in the Urine".
 Journal of Biological Chemistry 11 523 (1912)
- K.G.Falk and K. Sugiera
 "A Comparative Study of Aeration and Heat
 Distillation in the Kjeldahl Method for the
 Determination of Nitrogen".
 Journal of the American Chemical Society
 38 916 (1916)
- M. Nencki and J. Zaleski
 "The Determination of Ammonia in Animal Fluids
 and Tissue".
 Archiv für Experimentelle Pathologie
 36 18 (1895)
- Alfred H. Allen
 "Error of Hypobromite Method for Urea".
 Commercial Organic Analysis Vol. III, Part III 270.
- Soldern
 "Determination of Ammonia".
 Zeitschrift für Biologie 38 236 (1899)

RESPIRATION.

- A. Leimdorfer
 "The Conditions of Gas Tensions in the Lungs
 which Stimulate Respiration".
 Biochemie Zeitschrift 22 45 (1910)
- August Krogh
 "The Mechanism of the Gas Exchange in the Lungs".
 Skandinavisches Archiv für Physiologie (1910)
- L.E.Emmes and J.A.Riche
 "The Respiratory Exchange as Affected by Body
 Position".
 American Journal of Physiologie 138 167 (1911)
- Hans Winterstein
 "Regulation of Respiration by the Blood".
 Archiv der gesamten Physiologie 138 167 (1911)
- J. Lindhard
 "On the Excitability of the Respiratory Centre".
 Journal of Physiology 42 337 (1911)
- K.A.Hasselbalch
 "Regulation of Neutrality and the Sensitiveness
 of the Respiratory Centers in its Action on the
 Tension of Carbon Dioxide of the Blood".
 Biochemie Zeitschrift 46 405 (1913)

RESPIRATION (continued)

- J.M.H.Campbell, C.G.Douglas, J.S.Haldane and I.G.Hobson.
 "The Response of the Respiratory Center to
 Carbonic Acid, Oxygen and Hydrogen ion Concen-
 tration".
 Journal of Physiology 46 301 (1913)
- A.L.Sample and O.Porges
 "Relation of Carbon Dioxide Tension of the Blood
 to the Respiration".
 Biochemie Zeitschrift 54 182 (1914)
- J.M.H.Campbell, C.G.Douglas and F.G.Hobson
 "The Sensitiveness of the Respiratory Center to
 Carbonic Acid and Dead Space during Hyperpnea".
 Journal of Physiologie 48 303 (1914)
- Yandell Henderson
 "Variations in the Dead Space in Breathing",
 American Journal of Physiology 38 1 (1915)
- Hans Winterstein
 "Physico-chemical Regulation of Respiration."
 Biochemie Zeitschrift 70 45 (1915)
- H.L.Higgins and J.H.Means
 "Effects of Certain Drugs on the Respiration
 and Gaseous Metabolism in Normal Human Subjects".
 Journal of Pharmacology and Experimental Thera-
 peutics". 7 1 (1915)
- J.S.Haldane
 "Variations in the Effective Dead Space in
 Breathing".
 American Journal of Physiology 38 20 (1915)
- F.W.Peabody
 "Some Aspects of the Clinical Study of the Res-
 piration".
 American Journal of Medical Science 151 184 (1916)
- A. Krogh and J. Lindhard
 "The Volume of the Dead Space in Breathing and
 the Mixing of Gases in the Lungs of Man".
 Journal of Physiology 51 59 (1917)
- K.R.Hooker, D.W.Wilson and Helene Connett
 "The Effect of Carbon Dioxide and Other Substances
 in the Respiratory and Cardio-vascular Centers".
 American Journal of Physiology 43 351 (1917)
- J.S.Haldane and J.G.Priestley
 "The Regulation of the Lung Ventilation".
 Journal of Physiology 32 225 (1905)
- Y. Henderson and H.W.Haggard
 "Respiratory Regulation of the Carbon Dioxide
 Capacity of the Blood".

RESPIRATION (continued)

- W.M.H. Campbell, C.G. Douglas, J.S. Haldane and I.C. Hobson.
"The response of the Respiratory Center to
Carbonic Acid, Oxygen and Hydrogen Ion Concen-
tration".
Journal of Physiology 46 301 (1913)
- A.L. Remond and G. Borge.
"Relation of Carbon Dioxide Tension of the Blood
to the Respiration".
Biochimica et Biophysica Acta 182 182 (1914)
- W.M.H. Campbell, C.G. Douglas and I.C. Hobson.
"The sensitivity of the Respiratory Center to
Carbonic Acid and Ventilation during Hypoxia".
Journal of Physiology 48 303 (1914)
- Yandell Henderson.
"Variations in the Dead Space in Breathing".
American Journal of Physiology 1 38 (1915)
- Hans Hildebrandt.
"Physico-chemical Regulation of Respiration".
Biochimica et Biophysica Acta 48 48 (1915)
- B.L. Higgins and J.S. Haldane.
"Effects of Certain Drugs on the Respiration
and Gaseous Metabolism in Normal Human Subjects".
Journal of Pharmacology and Experimental Therapies-
1 1 (1915)
- J.S. Haldane.
"Variations in the Effective Dead Space in
Breathing".
American Journal of Physiology 38 30 (1915)
- F.W. Peabody.
"Some Aspects of the Clinical Study of the Res-
piration".
American Journal of Medical Science 151 184 (1915)
- A. Krogh and J. Lindhard.
"The Volume of the Dead Space in Breathing and
the Mixing of Gases in the Lungs of Man".
Journal of Physiology 59 51 (1915)
- E.B. Hooker, J.W. Wilson and Haldane Comfort.
"The Effect of Carbon Dioxide and Other Substances
in the Respiratory and Cardio-vascular Centers".
American Journal of Physiology 43 381 (1917)
- J.S. Haldane and J.G. Priestley.
"The Regulation of the Lung Ventilation".
Journal of Physiology 58 323 (1905)
- Y. Henderson and R.W. Haggard.
"Respiratory Regulation of the Carbon Dioxide
Capacity of the Blood".

RESPIRATION (continued)

Motiv Yamada

"The Haldane-Henderson Method for Estimating the Tension of Alveolar Carbon Dioxide and the Influence of Oxygen on the Stimulability of the Respiratory Centers".

A. J. Journal of the Chemical Society 114 I 511 (1918)

Eugen Jenni

"Estimations of the Tension of Carbon Dioxide in the Alveolar Air by the Henderson-Russel Modification of Haldane's Method and their Application for Testing the Stimulability of the Breathing Centre".

Journal of the Chemical Society 114 II 1462 (1918)

R.G.Pearce

"The Physiology of Respiration".

American Journal of Physiology 44 369 (1918)

L.G.Benedict and T.M.Carpenter

"Respiration Colorimeter for Studying the Respiratory Exchange of Man".

Publ. Carnegie Institute of Washington (1913)

J.S.Haldane

"Organism and Environment as Illustrated by the Physiology of Breathing".

Yale University Press (1917)

J.P.Peters, Jr.

"The Response of the Respiratory Mechanism to Rapid Changes in the Reaction of the Blood".

American Journal of Physiology 44 84 (1918)

H.W.Haggard and Y.Henderson

"Haemato-respiratory Function".

Journal of Biological Chemistry 39 163 (1919)

H. Hirst, J. Beckman, H. Hirst and H. Netterlester
"Alveolar Air Analysis".

Deutscher Archiv für Klinische Medizin

117 304-308 (1918)

F.S. Barker and C. Frothingham

"A Comparison in Various Methods of the Carbon Dioxide Tension in the Alveolar Air with the Amount of Carbon Dioxide in the Venous Blood".

Archives of Internal Medicine 18 306 (1918)

ALVEOLAR CARBON DIOXIDE.

Theodore Hough

"The Influence of Muscular Activity upon the
Alveolar Tensions of Oxygen and Carbon Dioxide".
American Journal of Physiology 30 18 (1912)

O.Porges, A.Leimdorfer and E.Markovici

"The Carbon Dioxide Tension of the Blood in
Pathological Conditions".
Zeitschrift klinische Medizin 73 389 (1912)

A.Krogh and J. Lindhard

"The Average Composition of the Alveolar Air and
Its Variations during the Respiratory Cycle".
Journal of Physiology 47 431 (1914)

H.L.Higgins

"The Influence of Food, Posture and Other Factors
on the Alveolar Carbon Dioxide Tension in Man".
American Journal of Physiology 34 114 (1914)

L.S.Fridericia

"The Determination of Diabetic Acidosis by a
Study of the Carbon Dioxide Tension of the
Alveolar Air".
Zeitschrift für klinische Medizin 80 1 (1914)

E.P.Poulton

"Observation on Carbon Dioxide in Alveolar Air of
Diabetes in Relation to the Onset of Coma".
Proceedings of the Royal Society of Medicine
7 No.7 171 (1915)

Paul Roth

"Estimations of Carbon Dioxide in Alveolar Air".
Journal of the American Medical Association.
65 413 (1915)

J.Howland and W.McK. Marriott

"Observations upon the So-called Food Intoxica-
tions of Infants with Special Reference to the
Alveolar Air".
Proceedings of the Society for Experimental
Biology and Medicine 12 51 (1915)

H.Straub, K.Beckman, H.Erdt and M. Mettenlecker

"Alveolar Air Analyses".
Deutscher Archiv für klinische Medizin
117 304-539 (1915)

I.C.Walker and C.Frothingham

"A Comparison in Various Diseases of the Carbon
Dioxide Tension in the Alveolar Air with the
Amount of Carbon Dioxide in the Venous Blood".
Archives of Internal Medicine 18 304 (1916)

ALVEOLAR CARBON DIOXIDE.

- Theodore Hough
"The Influence of Muscular Activity upon the
Alveolar Tension of Oxygen and Carbon Dioxide".
American Journal of Physiology 30 18 (1912)
- O. Forster, A. Latschewer and W. Markovitch
"The Carbon Dioxide Tension of the Blood in
Pathological Conditions".
Zeitschrift für Klinische Medizin 78 282 (1912)
- A. Krogh and J. Lindhard
"The Average Composition of the Alveolar Air and
Its Variations during the Respiratory Cycle".
Journal of Physiology 44 431 (1914)
- E. I. Higgins
"The Influence of Food, Exercise and Other Factors
on the Alveolar Carbon Dioxide Tension in Man".
American Journal of Physiology 34 114 (1914)
- E. E. Riddle
"The Determination of Metabolic Acids by a
Study of the Carbon Dioxide Tension of the
Alveolar Air".
Zeitschrift für Klinische Medizin 80 1 (1914)
- E. E. Forster
"Observations on Carbon Dioxide in Alveolar Air of
Diabetics in Relation to the Gust of Comm".
Proceedings of the Royal Society of Medicine
Vol. 7 191 (1912)
- Paul Roth
"Observations of Carbon Dioxide in Alveolar Air".
Journal of the American Medical Association 35 412 (1915)
- J. Howell and W. Hock, Marriot
"Observations upon the so-called Food Intoxica-
tions of Patients with Special Reference to the
Alveolar Air".
Proceedings of the Society for Experimental
Biology and Medicine 12 51 (1915)
- H. Girard, V. Beckman, E. Hirt and W. Mettenleiter
"Alveolar Air Analyses".
Deutscher Archiv für Klinische Medizin 114 304-333 (1915)
- I. C. Walker and C. Frothingham
"A Comparison in Various Diseases of the Carbon
Dioxide Tension in the Alveolar Air with the
Amount of Carbon Dioxide in the Venous Blood".
Archives of Internal Medicine 18 304 (1915)

ALVEOLAR CARBON DIOXIDE (continued)

- E.D.Friedman and H.C.Jackson
"The Carbon Dioxide Content of the Blood
and Alveolar Air in Obstructed Respiration".
Proceedings of the Society for Experimental
Biology and Medicine. 13 73 (1917)
- D.D.Van Slyke, E.Stillman and G.E.Cullen.
"Alveolar Carbon Dioxide and Plasma Bicarbonate
in Normal Men during Digestive Rest and Activity."
Journal of Biological Chemistry 31 217 (1917)
- Yandell Henderson and W.H.Morriss
"The Determination of Carbon Dioxide in the
Alveolar Air and Blood and the Carbon Dioxide
Combining Power of the Plasma and the Whole
Blood".
Journal of Biological Chemistry 31 217-27 (1917)
- R.G.Pearce
"The Capacity of the Air Passages and the
Percentage of Carbon Dioxide in Alveolar Air
during Rest and Exercise".
American Journal of Physiology 43 73 (1917)
- Yandell Henderson and A.L.Prince
"The Carbon Dioxide Tension of the Venous Blood
and the Circulation Rate".
Journal of Biological Chemistry 32 325 (1917)
- C.Sonne
"Arterial and Venous Carbon Dioxide Tensions".
Deutscher Archiv für klinische Medizin
124 358 (1918)
- L.S.Fridericia
"The Oxygen and Carbon Dioxide Tension in the
Blood of the Pulmonary Artery".
Biochemie Zeitschrift 85 307 (1918)
- H.Laurens
"Effect of Holding the Breath and of Rebreathing
on the Carbon Dioxide Tensions".
American Journal of Physiology 46 147 (1919)
- O.M.Schlos and H.Harrington
"Comparison of the Carbon Dioxide Tension of the
Alveolar Air and the Hydrogen ion Concentration
of the Urine with the Bicarbonate of Blood
Plasma".
American Journal of Diseases of Children
17 85 (1919)

ALCOHOLIC CARBON DIOXIDE (continued)

H.C. Williams and E.C. Jackson
"The Carbon Dioxide Content of the Blood
and Alveolar Air in Obstructed Respiration."
Proceedings of the Society for Experimental
Medicine and Biology. 13 73 (1917)

C.D. Van Slyke, H. Williams and G.H. Gilson
"Alveolar Carbon Dioxide and Plasma Bicarbonate
in Normal Men during Obstructive Rest and Activity."
Journal of Biological Chemistry 31 217 (1917)

Yandell Henderson and W.H. Norris
"The Determination of Carbon Dioxide in the
Alveolar Air and Blood and the Carbon Dioxide
Combining Power of the Plasma and the Whole
Blood."
Journal of Biological Chemistry 31 217-227 (1917)

H.C. Berge
"The Capacity of the Air Passages and the
Percentage of Carbon Dioxide in Alveolar Air
during Rest and Exercise."
American Journal of Physiology 43 78 (1917)

Yandell Henderson and A.E. Prince
"The Carbon Dioxide Tension of the Venous Blood
and the Circulation Rate."
Journal of Biological Chemistry 32 325 (1917)

G. Sonne
"Arterial and Venous Carbon Dioxide Tensions."
Doklady Akad. Nauk SSSR 124 388 (1918)

I.E. Wigglesworth
"The Oxygen and Carbon Dioxide Content in the
Blood of the Pulmonary Artery."
Biochimica Biophysica Acta 35 307 (1918)

H. Tammann
"Effect of Holding the Breath and of Rebreathing
on the Carbon Dioxide Tension."
American Journal of Physiology 45 147 (1918)

G.H. Scholten and H. Harrington
"Comparison of the Carbon Dioxide Tension of the
Alveolar Air and the Hydrogen Ion Concentration
of the Urine with the Bicarbonate of Blood
Plasma."
American Journal of Diseases of Children 17 85 (1919)

ALVEOLAR CARBON DIOXIDE (continued)

G.A.Harrop, Jr.

"Oxygen and Carbon Dioxide Content of the Arterial and Venous Blood in Normal Individuals and in Patients with Anaemia and Heart Disease".

Journal of Experimental Medicine 30 241 (1919)

ALVEOLAR CARBON DIOXIDE Methods of Determination.

J.S.Haldane

"Methods of Air Analysis".

Published Charles Griffin and Co. London. (1912)

Yandell Henderson and W.G.Russell

"A Simple Method for the Determination of the Carbon Dioxide Content of the Alveolar Air by Means of Baryta".

American Journal of Physiology 29 436 (1912)

M. Hahn and R. Heim

"The Determination of Carbon Dioxide in Alveolar Air by means of the Interferometer".

Berliner klinische Wochenschrift 50 197 (1913)

L.S.Fridericia

"The Determination of Diabetic Acidosis by means of the Carbon Dioxide in the expired Air".

Munchener Medizinische Wochenschrift 60 2358 (1914)

W.M.Boothby and F.W.Peabody

"A Comparison of Methods for Obtaining Alveolar Air".

Archives of Internal Medicine 13 497 (1914)

T.M.Carpenter

"A Comparison of Methods for Determining the Respiratory Exchange of Man".

Carnegie Institute Publication 216 (1915)

W.McK. Marriott

"Determination of Alveolar Carbon Dioxide Tensions by a Simple Method".

Journal of the American Medical Association 66 1594 (1916)

J.R.Williams

"Clinical Methods of Measuring Acidosis".

Medical Record 90 223 (1916)

Sir James Dewar

"Diffusion of Gases through India Rubber".

Journal of the Chemical Society 114 II 186 (1918)

ALVEOLAR CARBON DIOXIDE (continued)

G.A. Hannon, Jr.
"Carbon and Carbon Dioxide Content of the
Arterial and Venous Blood in Normal Indi-
viduals and in Patients with Anemia and
Heart Disease".
Journal of Experimental Medicine 30 241 (1919)

ALVEOLAR CARBON DIOXIDE Methods of Determination.

L.S. Williams
"Methods of Air Analysis".
Published Charles Griffin and Co. London. (1912)

Yoshiji Hatanaka and E.C. Ross
"A Simple Method for the Determination of the
Carbon Dioxide Content of the Alveolar Air
by Means of a Syringe".
American Journal of Physiology 29 435 (1912)

H. Hahn and E. Bohr
"The Determination of Carbon Dioxide in Alveolar
Air by Means of the Interferometer".
Berliner Klinische Wochenschrift 86 197 (1913)

L.S. Williams
"The Determination of Alveolar Air by
Means of the Carbon Dioxide in the expired
Air".
Thomson Medical and Pharmaceutical 60 235 (1914)

H.M. Rosenthal and E.T. Rosenthal
"Comparison of Methods for Obtaining
Alveolar Air".
Archives of Internal Medicine 93 497 (1914)

T.H. Carpenter
"A Comparison of Methods for Determining the
Respiratory Exchange of Man".
Carnegie Institute Publication 216 (1915)

F.M.K. Hirst
"Determination of Alveolar Carbon Dioxide
by a Simple Method".
Journal of the American Medical Association 60 1534 (1916)

L.S. Williams
"Alveolar Methods of Measuring Ventilation".
Medical Record 90 225 (1916)

W. Loomis Brown
"Determination of Gases by a Simple Method".
Journal of the Chemical Society 111 155 (1918)

ALVEOLAR CARBON DIOXIDE

Methods for Determination (continued)

Pettersson

"Gas Analysis"

Zeitschrift für Analytische Chemie 25 467 (1886)

L.M.Denis

"Gas Analysis".

MacMillan Co.

(1913)

Walther Hempel

"Methods of Air Analysis",

Translated from 3d German Edition and

enlarged by L.M.Denis. MacMillan Co.

(1906)

Anonymous

"Carbon Dioxide Thermoscope"

Engineering

95 787

(1913)

Anonymous

"A Simple Carbon Dioxide Meter"

American Gas Light Journal

99

62

(1913)

Y.Henderson

"Haldane Gas Analyzer"

Journal of Biological Chemistry

38

(1918)

Augustus H. Gill

"Gas and Fuel Analysis for Engineers".

John Wiley and Sons, New York.

(1897)

Wilfred W. Scott

"Standard Methods of Chemical Analysis"

Second Edition. D. Van Nostrand Co., N.Y.

(1918)

Orsat

"Gas Analysis"

Chemical News.

29

177

(1874)

F.M.Seibert and W.C.Harpter

"Use of the Interferometer in Gas Analysis".

Bureau of Mines Technical Paper 185 18

(1918)

J.J.R.Macleod and R.L.Long

"Simplified Gas Analysis".

Journal of Laboratory and Clinical Medicine

4

564

(1919)

ANALYTICAL CHEMISTRY

Methods for Determination (continued)

- Patterson
"Gas Analysis"
Zeitschrift für Analytische Chemie 58 487 (1933)
- L.M. Denis
"Gas Analysis".
MacMillan Co. (1913)
- Walter Rappaport
"Methods of Air Analysis",
Translated from 25 German Editions and
enlarged by L.M. Denis. MacMillan Co. (1933)
- Anonymous
"Carbon Dioxide Thermostat"
Engineering (1913) 95 787
- Anonymous
"A Simple Carbon Dioxide Meter"
American Gas Light Journal (1913) 92 68
- Y. Hesterton
"Helium Gas Analyzer"
Journal of Biological Chemistry (1913) 38
- Augustus E. Gill
"Gas and Fuel Analysis for Engineers".
John Wiley and Sons, New York. (1927)
- Wilfred F. Scott
"Standard Methods of Chemical Analysis"
Second Edition. J. Van Nostrand Co., N.Y. (1913)
- Orsat
"Gas Analysis"
Chemical News. (1874) 33 177
- F.M. Beilby and F.C. Harper
"Use of the Interferometer in Gas Analysis".
Bureau of Mines Technical Paper 138 (1913)
- S.L.R. MacLeod and E.L. Long
"Simplified Gas Analysis".
Journal of Laboratory and Clinical Medicine (1913) 4 584

BOSTON UNIVERSITY



1 1719 02544 0944



ACCOPRESS BINDER

BF 250-P7 EMB

MADE BY

ACCO PRODUCTS, INC.

OGDENSBURG, N. Y.

